

NOTICE OF EXPRESS MAILING

Express Mail Mailing Label Number: EV326919981US

Date of Deposit with USPS: April 15, 2004

Person making Deposit: Christopher Haughton

APPLICATION FOR LETTERS PATENT

for

**NEWCASTLE DISEASE VIRUS INFECTIOUS CLONES, VACCINES AND  
NEW DIAGNOSTIC ASSAYS**

Inventors:

Bernardus P.H. Peeters  
Olav S. de Leeuw  
Guus Koch  
Arnoud L.J. Gielkens

Attorneys:  
Allen C. Turner  
Registration No. 33,041  
TRASKBRITT  
P.O. Box 2550  
Salt Lake City, Utah 84110  
(801) 532-1922

# **NEWCASTLE DISEASE VIRUS INFECTIOUS CLONES, VACCINES AND NEW DIAGNOSTIC ASSAYS**

## **CROSS-REFERENCE TO RELATED APPLICATIONS**

**[0001]** This application is a divisional application of U.S. Patent Application Serial No. 10/788,232, filed February 26, 2004, pending, which is a divisional application of U.S. Patent Application Serial No. 09/741,744, filed December 19, 2000, now U.S. Patent 6,719,979 B2, which application claims priority to and is the national phase of International Application No. PCT/NL99/00377, filed on 17 June 1999, designating the United States of America, the contents of which are incorporated by this reference.

## **TECHNICAL FIELD**

**[0002]** The invention relates generally to biotechnology, and more particularly to Newcastle disease virus ("NDV") infections of poultry.

## **BACKGROUND**

**[0003]** NDV is one of the most diverse and deadly avian pathogens. The almost simultaneous occurrence of Newcastle disease as an apparent new disease in several different geographical locations and the great variation in the severity of the disease has caused some problems with nomenclature.

**[0004]** The disease has been termed pseudo fowl pest, pseudo poultry plague, avian pest, avian distemper and avian pneumoencephalitis. The importance of the disease is primarily due to the development of the poultry industry during the 20th Century into a highly efficient international industry which is dependent on intensive trade between countries.

**[0005]** It is generally assumed that the first outbreaks of Newcastle disease occurred in 1926 in Java, Indonesia, and in Newcastle-upon-Tyne, England (Kraneveld, 1926; Doyle 1927). The name "Newcastle disease" was coined by Doyle as a temporary name to avoid a descriptive name that might be confused with other diseases. It later became clear that other less severe diseases were caused by viruses indistinguishable from NDV. In the US, a relatively mild

respiratory disease was named “avian pneumoencephalitis” and was shown to be caused by NDV (Beach, 1944). Within a few years, numerous NDV isolations that caused extremely mild or no disease in chickens were made around the world.

[0006] The following methods have been implicated in the spread of the disease: 1) movement of live birds, feral birds, game birds, racing pigeons and commercial poultry; 2) movement of people and equipment; 3) movement of poultry products; 4) airborne spread; 5) contaminated poultry feed; 6) contaminated water; 7) incompletely inactivated or heterogeneous vaccines. According to the OIE, Newcastle disease is a disease of poultry caused by a virus of avian-paramyxovirus serotype 1 (APMV-1) which has an intracerebral pathogenicity index (ICPI) in day-old chicks of 0.17 or greater. Virulent virus can also be confirmed by the presence of multiple basic amino acids at the C-terminus of the F2 protein and F (phenylalanine) at residue 117, the N-terminus of the F1 protein. Failure to demonstrate this amino acid sequence would require characterization by ICPI tests. The word “poultry” refers to domestic fowl, turkeys, guinea fowl, ducks, geese, quails, pigeons, pheasants, partridges and ratites that are reared or kept in captivity for breeding, the production of meat or eggs for consumption, or for restocking supplies of game.

[0007] According to Alexander (1988) three panzootics of Newcastle disease have occurred since the first recognition of the disease. The first represented the initial outbreaks of the disease and appears to have arisen in Southeast Asia. Isolated outbreaks, such as the one in England in 1926, were chance introductions ahead of the mainstream which slowly moved through Asia to Europe.

[0008] A second panzootic appears to have begun in the Middle East in the late 1960's and reached most countries by 1973. The more rapid spread of the second panzootic was probably caused by the major revolution of the poultry industry with considerable international trade.

[0009] A third panzootic primarily affected domesticated birds such as pigeons and doves (Vindevogel and Duchatel, 1988). The disease apparently arose in the Middle East in the late 1970's. By 1981, it reached Europe and then spread rapidly to all parts of the world, largely as a result of contact between birds at races and shows and the international trade in such birds.

[0010] Nowadays, Newcastle disease is still widespread in many countries of Asia, Africa, the Americas, and Europe. Only the countries of Oceania appear to be relatively free from the disease (Spradbrow, 1988).

[0011] NDV belongs to the order *Monomegavirales*, family *Paramyxoviridae*, subfamily *Paramyxoviridae*, and genus *Rubulavirus*. Apart from NDV, generally called avian-paramyxovirus type-1, eight other serotypes, designated avian-paramyxovirus type-2 to -9, can be distinguished on the basis of their antigenic relatedness in hemagglutination-inhibition tests and serum neutralization tests (Alexander, 1993).

[0012] Despite the consistency of the serological grouping there are some cross-relationships between viruses of the different serotypes.

[0013] The genome of NDV is a single-stranded RNA molecule of negative polarity, complementary to the messenger RNA's which code for the virus proteins. The RNA genome is approximately 15,200 nt in size and codes for the following gene products (listed from the 3' end to the 5' end of the genomic RNA): nucleocapsid protein (NP), phosphoprotein (P), matrix protein (M), fusion protein (F), hemagglutinin-neuraminidase (HN), and large polymerase protein (L) (Chambers et al., 1986).

[0014] The RNA is complexed with the NP, P and L proteins to form a ribonucleocapsid particle (RNP) that is surrounded by an envelope that is lined at the inside by the M protein. The envelope contains the F and HN proteins which are involved in attachment and penetration of the host cell.

[0015] Replication of NDV is similar to the strategy used by other paramyxovirinae. The initial step is attachment of the virus to the host cell receptors, mediated by the HN protein. Fusion of the viral envelope with the host cell membrane is dependent on the action of both the HN and F proteins and results in the release of the RNP into the cytoplasm where virus replication takes place.

[0016] The viral RNA-dependent RNA polymerase (which is part of the RNP) produces complementary transcripts that act as mRNA's and are used by the cell's translation machinery for the synthesis of virus proteins. Due to the accumulation of NP protein, the RNA

polymerase complex switches from transcription to replication, resulting in the synthesis of full-length genomic and antigenomic RNA molecules.

**[0017]** Newly formed RNP's are encapsidated at the cellular membrane by the action of the M protein and the F and HN proteins which have accumulated in the cellular plasma membrane. Newly formed virus particles are released from the infected cell by a budding mechanism. For more detailed information about NDV replication see Peeples (1988). For a recent review of the molecular biology of paramyxovirinae see Lamb and Kolakofsky (1996).

**[0018]** Apart from commercial domestic poultry (*e.g.*, chicken, turkey, pheasant, guinea fowl, duck, goose, and pigeon), a wide range of captive, semi-domestic and free-living birds, including migratory waterfowl, is susceptible to NDV and can be primary infection sources (Kaleta and Baldauf, 1988).

**[0019]** The pathogenicity of NDV strains differs greatly with the host. The most resistant species appear to be aquatic birds while the most susceptible are gregarious birds forming temporary or permanent flocks. Chickens are highly susceptible but ducks and geese may be infected and show few or no clinical signs, even with strains which are lethal for chickens.

**[0020]** Newcastle Disease is complicated in that different isolates and strains of the virus may induce enormous variation in the severity of the disease. Beard and Hanson (1984) grouped NDV strains and isolates into different pathotypes that relate to disease signs that may be seen in fully susceptible chickens: 1) viscerotropic velogenic NDV, which produces acute lethal infections in which hemorrhagic lesions are prominent in the gut; and neurotropic velogenic NDV, which produces high mortality preceded by respiratory and neurological signs, but no gut lesions; 2) mesogenic NDV, which produces low mortality, acute respiratory disease and nervous signs in some birds; 3) lentogenic NDV, which produces mild or unapparent respiratory infections or even asymptomatic enteric NDV, avirulent viruses that appear to replicate primarily in the intestinal tract. Some overlap between the signs associated with the different groups has been reported.

**[0021]** The virus enters the body via the respiratory and the intestinal tract or via the eye. In the trachea, the virus is spread by ciliary action and by cell-to-cell spread. After initial

multiplication at the introduction site, virus is carried during episodes of viremia to spleen, liver, kidney and lungs. Viruses of some strains reach vital organs like liver and kidney very rapidly so that the birds may die before disease symptoms are overt.

**[0022]** Most viruses reach the central nervous system via the blood before significant amounts of antibody exist. A long, asymptomatic carrier state presumed to occur in psittacines constitutes a potential threat to the poultry industry. A long term carrier state of both lentogenic and velogenic virus may also exist in chickens (Heuschele and Easterday, 1970).

**[0023]** During the replication of NDV it is necessary for the precursor glycoprotein Fo to be cleaved to F1 and F2 for the progeny virus to be infectious (Rott and Klenk, 1988). This posttranslational cleavage is mediated by host cell proteases. If cleavage fails to take place, non-infectious virus particles are produced and viral replication cannot proceed. The Fo protein of virulent viruses can be cleaved by a wide range of proteases, but Fo proteins in viruses of low virulence are restricted in their sensitivity and these viruses can only grow *in vivo* in certain host cell types and in general cannot be grown *in vitro*.

**[0024]** Lentogenic viruses only replicate in areas with trypsin-like enzymes such as the respiratory and intestinal tract, whereas virulent viruses can replicate in a range of tissues and organs resulting in fatal systemic infection.

**[0025]** Amino acid sequencing of the Fo precursor has shown that low-virulence viruses have a single arginine (R) that links the F2 and F1 chains, whereas virulent strains possess additional basic amino acids forming two pairs such as K/R-X-K/R-R-F at the site of cleavage. Furthermore, the F2 chain of virulent strains generally starts with a phenylalanine residue whereas that of non-virulent strains generally starts with a leucine.

**[0026]** For a few strains of NDV the HN protein is also produced as a precursor that requires cleavage to be biologically active (Garten et al., 1980; Millar et al., 1988).

**[0027]** Besides cleavability of the F and HN proteins, other viral factors may contribute to pathogenicity. Madansky and Bratt (1978, 1981a, 1981b) have shown that alterations in transcription and translation could modulate growth and cell-to-cell spread of the virus and/or cytopathogenicity.

**[0028]** The initial immune response to infection with NDV is cell mediated and may be detectable as early as 2-3 days after infection with live vaccine strains. This presumably explains the early protection against challenge that has been recorded in vaccinated birds before a measurable antibody response is seen (Gough and Alexander, 1973).

**[0029]** At about 1 week after infection, circulating antibodies may protect the host from re-infection. In the early phase IgM is involved, followed by IgG. Titers and protection peak after about 3 weeks, and gradually decline if without boosting. This means that for older birds, re-vaccinations are necessary.

**[0030]** Only live vaccines administered by the respiratory route stimulate antibody in all mucosal surfaces as well as in serum. Inactivated vaccine, even when applied via the intramuscular route, does not elicit local resistance in the respiratory tract, despite high concentrations of serum antibody.

**[0031]** This stresses the importance of live vaccines capable of presenting viral antigen to the upper respiratory tract to induce both local and systemic immunity. Small droplets penetrate into the lower respiratory tract thereby provoking a mainly humoral immune response, while coarse droplets stimulate local immunity in the upper respiratory tract.

**[0032]** Therefore, aerosols with a wide range of droplet sizes generate the best overall local and humoral immunity.

**[0033]** It should be noted, however, that despite intensive vaccination with current vaccines creating high levels of antibody titers, virus can still be recovered from mucous surfaces.

**[0034]** The identification of Newcastle disease in the USA led to the use of inactivated vaccines (Hofstad, 1953). The observation that some of the enzootic viruses produced only mild disease resulted first in the development of the mesogenic live vaccine Roakin (Beaudette et al., 1949) and, subsequently, in the development of the milder Hitchner B1 (Hitchner and Johnson, 1948) and LaSota (Goldhaft, 1980) strains, which are now the most widely used live vaccines.

**[0035]** NDV live vaccines can be divided into two groups, lentogenic and mesogenic. Mesogenic strains are suitable only for secondary vaccination of birds due to their greater virulence. The immune response increases as the pathogenicity of the live vaccine increases.

Therefore, to obtain the desired level of protection without serious reaction, currently vaccination programs are used that involve sequential use of progressively more virulent vaccines, or live vaccines followed by inactivated vaccines.

**[0036]** One of the main advantages of live vaccines is that they may be administered by inexpensive mass application techniques. A common method of application is via drinking water. However, drinking water application must be carefully monitored as the virus may be inactivated by excessive heat and light and by virucidal impurities in the water.

**[0037]** Mass application of live vaccines by sprays and aerosols is also very popular due to the ease with which large numbers of birds can be vaccinated in a short time. It is important to achieve the correct particle size by controlling the conditions under which the particles are generated.

**[0038]** Currently used live vaccines have several disadvantages. The vaccine may still cause disease signs, depending upon environmental conditions and the presence of complicating infections. Therefore, it is important to use extremely mild virus for primary vaccination and, as a result, multiple vaccinations are usually needed. Furthermore, maternally derived antibodies may prevent successful primary vaccination with lentogenic live vaccines.

**[0039]** Inactivated vaccines are usually produced from infectious allantoic fluid which is treated with formalin or beta-propiolactone to kill the virus and mixed with a suitable adjuvant. Inactivated vaccines are administered by injection, either intramuscularly or subcutaneously. Inactivated vaccines are expensive to produce and to apply.

**[0040]** However, inactivated oil-emulsion vaccines are not as adversely affected by maternal immunity as live vaccines and they can be used in day-old chicks. Advantages of inactivated vaccines are the low level of adverse reactions in vaccinated birds, the high level of protective antibodies, and the long duration of protection. None of the above vaccines can serologically be differentiated from wild-type NDV.

**[0041]** The development of recombinant viral vaccines has been of interest to the poultry industry for a number of years. The concept is to insert genes of critical immunizing epitopes of a disease agent of interest into a nonessential gene of a vector virus. Vaccination



with the recombinant virus thus results in immunization against both the vector virus as well as the disease agent of interest.

[0042] Several types of viruses have been evaluated as potential live viral vaccines for poultry. Two avian viruses that have received most attention are fowl pox virus (FPV) and herpes virus of turkeys (HVT). Fowl pox virus is a DNA virus that has a large genome and hence is considered to have ample room to carry foreign genes.

[0043] When attenuated, FPV does not cause clinical disease and is commonly used as a vaccine in chickens. HVT is also a DNA virus and is classified as serotype III of the Marek's disease virus (MDV) family. HVT is non-pathogenic for chickens yet cross-protective against MDV and is commonly used to vaccinate chickens against Marek's disease.

[0044] It has been shown that protection against Newcastle disease can be induced by using recombinant HVT or FPV vaccines (Morgan et al., 1992, 1993; Heckert et al., 1996; Boursnell et al., 1990; Taylor et al., 1990).

[0045] However, the onset of protection against Newcastle disease following vaccination with such recombinant vaccines that express either the NDV F protein or both the F and HN proteins was severely delayed compared to that following vaccination with a conventional live or inactivated NDV vaccine, possibly because the recombinant vaccines do not provide a wide enough immunological specter of antigenically relevant NDV epitopes other than those found on the NDV protein that is expressed by the recombinant vaccine or are not properly presented to the immune system.

[0046] Furthermore, local (mucosal, respiratory or enteric) protection was not effectively induced in birds vaccinated with the recombinants. This is a serious drawback since vaccines used for primary vaccination against respiratory diseases must induce local immunity to prevent infection and spread of virulent viruses that infect chickens reared under field conditions.

[0047] Antibodies against NDV which are capable of protecting the host can be measured in virus neutralization tests. However, since the neutralization response appears to parallel the hemagglutination-inhibition (HI) response, the latter test is frequently used to assess the protective response, especially after vaccination.

[0048] Antibodies against both the F and HN proteins can neutralize NDV. However, antibodies against the F protein appear to induce greater neutralization than those directed against HN in *in vivo* and *in vitro* tests (Meulemans et al., 1986).

[0049] The presence of specific antibodies to NDV in the serum of a bird gives little information on the infecting strain of NDV and therefore has limited diagnostic value.

[0050] The omnipresence of lentogenic NDV strains in birds in most countries and the almost universal use of live vaccines that cannot be distinguished, at least not serologically from wild-type NDV, mean that the mere demonstration of infection is rarely adequate cause for control measures to be imposed. Since field disease may be an unreliable measure of the true virulence of the virus, it is necessary to further characterize the virus that is found.

[0051] At present, the only method of Newcastle disease diagnosis which allows characterization of the infecting strain is virus isolation followed by pathogenicity testing. At present, three *in vivo* tests are used for this purpose: 1) mean death time (MDT) in eggs; 2) intracerebral pathogenicity index (ICPI) in one-day-old chickens; 3) Intravenous pathogenicity index (IVPI) in 6-week-old birds.

[0052] These tests suffer from a number of drawbacks, such as the availability of animals, poor reproducibility, and the relatively long duration of the tests. Last but not least, these tests do not allow a simple serological identification of poultry vaccinated with a vaccine or infected with a wild-type strain.

[0053] As an alternative to *in vivo* tests, the polymerase chain reaction (PCR) has been successfully used to distinguish between virulent and non-virulent and non-virulent isolates (Stauber et al., 1995; Kant et al., 1997), however, again serological differentiation is not possible.

[0054] The raising of poultry and trade of their products is now organized on an international basis, frequently under management of multinational companies. The threat of Newcastle disease has proven a great restraint on such trade.

[0055] Successful control of Newcastle disease will only be approached when all countries report outbreaks. However, international agreements are not simple due to enormous variation in the extent of disease surveillance in different countries. Some countries do not

vaccinate and would not want any form of NDV introduced in domestic poultry because vaccinated poultry cannot be distinguished from those infected with wild-type NDV.

[0056] Others only allow the use of specific live vaccines and consider other vaccines as unacceptably virulent. Yet other countries have the continued presence of circulating highly virulent virus, which is not recognized as such because overt disease is masked by vaccination.

[0057] In many countries legislation exists to control Newcastle disease outbreaks that may occur. National control measures are directed at prevention of introduction and spread. Most countries have restrictions on trade in poultry products, eggs, and live poultry. Most countries have established quarantine procedures for importation, especially for psittacine birds.

[0058] Some countries have adopted eradication policies with compulsory slaughter of infected birds, their contacts, and products. Others require prophylactic vaccination of birds even in the absence of outbreaks, while some have a policy of ring vaccination around outbreaks to establish a buffer zone.

[0059] A need exists for better vaccines and for better diagnostic methods which can be used to control Newcastle disease. Due both to large differences in the dose that is received by individual birds during mass application of live vaccines and to variation in levels of maternal immunity in young chickens, post-vaccination reactions with live vaccines are inevitable. This is one of the main concerns of farmers in countries where vaccination is compulsory.

[0060] Furthermore, many vaccines are mixtures of sub-populations. When cloned, these sub-populations may differ significantly from each other in immunogenicity and pathogenicity (Hanson, 1988).

[0061] However, the largest drawback of currently used live vaccines and inactivated vaccines is the fact that vaccinated animals cannot be distinguished from infected animals with currently used screening techniques such as hemagglutination-inhibition or virus neutralization tests.

[0062] Virulent field-virus may still spread in vaccinated flocks since disease symptoms are masked by vaccination. Since virus isolation and characterization of virulence by *in vivo* techniques is not feasible on a large scale, there is a great need for new and effective attenuated live vaccines which can be serologically discriminated from field-viruses.

[0063] Such vaccines, called NDV marker vaccines (and accompanying diagnostic methods and kits) which should provide the fullest possible immunological specter of antigenically relevant NDV epitopes, and yet should be serologically distinct from wild-type NDV are not yet available.

#### DISCLOSURE OF THE INVENTION

[0064] The invention provides a method of modifying an avian-paramyxovirus genome by genetic modification. It also provides genetically modified avian paramyxovirus and an avian-paramyxovirus marker vaccine.

[0065] The advent of modern molecular biological techniques has allowed the genetic modification of many RNA viruses, including negative-strand RNA viruses. This technique is often referred to as “reverse genetics.” One first provides a (full-length) cDNA copy of the viral RNA, after which one transcribes this DNA in susceptible cells to produce infectious RNA which can again replicate to produce infectious virus particles.

[0066] In general, by previous modification of the cDNA with standard molecular biological techniques, it is possible to obtain a genetically modified RNA virus. However, this has never materialized for NDV or other avian paramyxoviruses, it has even not yet been possible to generate minigenome fragments or plasmids of avian-paramyxovirus genomic fragments to study replicative events of avian paramyxovirus, thereby creating an understanding on how to construct infectious copy virus.

[0067] Surprisingly, although in this description it has now been fully established that the genome of avian paramyxovirus is the smallest of all paramyxovirus genomes sequenced up to now, especially the 5'-terminal end sequence of the NDV genome is much longer than previously had been established and was expected by comparison with other *Paramyxoviridae*. The invention now for the first time provides a full sequence of an avian-Paramyxovirus genome and provides full-length or mini-genomic length cDNA of such a virus.

[0068] The invention herewith provides avian-paramyxovirus cDNA at least comprising a nucleic acid sequence corresponding to the 5'-terminal end of the genome of avian paramyxovirus allowing generating an infectious copy of avian paramyxovirus, the cDNA

preferably comprising a full-length cDNA. However, the invention also provides cDNA at least comprising a nucleic acid sequence corresponding to the 5'-terminal end of the genome of avian paramyxovirus thereby allowing generating a replicating avian-paramyxovirus minigenome. Such mini-genomes can advantageously be used to transcribe RNA and/or express protein from modified nucleic acid sequences. The invention provides a cDNA according to the invention at least partly derived from NDV, for example wherein the NDV is a lentogenic virus, preferably derived from a vaccine strain, such as LaSota strain ATCC VR-699.

**[0069]** The invention furthermore provides a cDNA according to the invention additionally provided with a modification, such as a deletion, insertion, mutation, reversion, or otherwise in a nucleic acid. For example, a cDNA is provided wherein the modification comprises a nucleic acid encoding a modified protease cleavage site, for example wherein the cleavage site is a protease cleavage site of the fusion (F) protein.

**[0070]** In yet another embodiment, the invention provides a cDNA according to the invention wherein the modification comprises a nucleic acid encoding a hybrid viral protein, such as a hybrid hemagglutinin-neuraminidase (HN) protein as described in the experimental part of the invention. The invention also provides a cDNA according to the invention wherein the modification comprises a deletion in a nucleic acid encoding a viral protein, such as a matrix (M) protein.

**[0071]** The invention additionally provides a cDNA according to the invention additionally provided with a nucleic acid encoding a heterologous antigen, preferably wherein the antigen is derived from a poultry pathogen, as for example described below. An RNA, and protein derived thereof, obtained from a cDNA according to the invention is also provided.

**[0072]** In recent years, a number of non-segmented negative-strand RNA viruses has been fully characterized and fundamental work on the replication and expression of their genomes has culminated in the ability to generate infectious virus entirely by transfecting cells with cloned cDNA of the virus (reviewed by Conzelmann, 1996).

**[0073]** To date, infectious virus of non-segmented negative-strand RNA viruses has been generated from cloned cDNA of for example rabies virus (Schnell et al., 1994, Conzelmann; EP0702085A1), vesicular stomatitis virus (Lawson et al., 1995; Whelan et al.,

1995), Sendai virus (Garcin et al., 1995), measles virus (Radecke et al., 1995; Schneider et al., 1997; EP0780475A1), human respiratory syncytial virus (Collins et al., 1995), rinderpest virus (Baron and Barrett, 1997), and human parainfluenza virus type 3 (Hoffman and Banerjee, 1997, Conzelmann; EP0702085A1), (Schnell et al., 1994; EP0702085A1).

**[0074]** However, all of above infectious copy viruses are capable of growing both *in vivo* as well as *in vitro* in hosts, tissues or cells of various origin, allowing easy cDNA transfection and replication and generation of infectious virus particles on a suitable cell line.

**[0075]** Such a possibility does not exist for NDV certainly not for lentogenic NDV strains which can provide a vaccine. Virulence of such an NDV strain is associated with its ability to replicate in a wide range of cells, reflected by the fact that virulent strains can easily replicate *in vitro* and *in vivo*, whereas vaccine strains can only replicate *in vivo*.

**[0076]** Thus, with NDV a “Catch 22” situation is apparent. While attempts to generate an infectious copy virus from for example infectious cDNA may possibly result in infectious virus, such virus is in general not suitable for use as a vaccine because the thus generated infectious virus is by default too virulent to be used as vaccine; the fact that it can be generated and replicated after transfection of cDNA on a cell line reflects its easy cleavability of the Fo protein into F1 and F2, as discussed above a hallmark of virulence of a NDV.

**[0077]** Using a vaccine strain as parent material for the cDNA would not solve this problem; a vaccine strain, especially of a lentogenic type does not contain an easily cleavable Fo protein, rendering it impossible for first generation virus to continue to replicate. The cell used for transfection will simply not be susceptible to support one or more rounds of replication of vaccine-type virus with a non-cleaved Fo protein.

**[0078]** The invention now elegantly provides a solution for this problem, and therewith provides infectious copy NDV, for example for use in a vaccine.

**[0079]** The invention provides a method to generate infectious copy NDV comprising transfecting cells, capable of expressing viral NP, P and L proteins for complexing with viral RNA with cloned full-length or genomic-length cDNA of the virus and further comprising incubating the cells in growth medium comprising proteolytic activity allowing cleavage of the Fo protein of the virus.

**[0080]** In our system, co-transfection of a plasmid-expressing NP could be omitted. NP is probably expressed from the full length cDNA because the NP gene is the first gene after the 5' end of the antigenomic RNA. Since eukaryotic mRNA are usually monocistronic, expression of distal genes is not expected. However it is possible to generate full-length cDNA in which the relative positions of the NDV genes are changed. If the first gene of such a cDNA is the P or L gene, it is not necessary to express the corresponding gene product from a co-transfected plasmid.

**[0081]** As an alternative to using full-length cDNA, it is possible to use two or more subgenomic cDNA's which generate replication competent subgenomic cDNA's and which together express the full complement of avian-paramyxovirus proteins. Even if the RNA's are packaged separately, the resulting virus-like particles can be used for successive rounds of replication by means of co-infection and complementation of gene functions.

**[0082]** In a preferred embodiment, the invention provides a method wherein the proteolytic activity is derived of an enzyme, such as a trypsin-like enzyme, or is derived of a composition comprising the proteolytic activity. In a much preferred embodiment, the growth medium comprises allantoic fluid comprising proteolytic activity. Cleavage of the Fo protein is required for the generation of infectious virus. It is possible to generate infectious virus from lentogenic strain without the addition of exogenous proteolytic activity. By inoculating the supernatant of transfected cells into the allantoic cavity of embryonated eggs, the proteolytic activity which is present in the allantoic fluid is able to cleave the Fo protein to generate the fusion-competent F1-F2 complex. Virions with such an activated F protein are able to infect susceptible cells and replication in cells which express the desired proteolytic activity yields infectious progeny. As an alternative to providing the desired proteolytic activity to the supernatant of transfected cells, it is, for example, possible to use a cell that is permissive for NDV and which already expresses the proteolytic activity. Such a cell line is used to produce infectious lentogenic NDV without the addition of exogenous proteolytic activity. Such a cell line can also be generated by stable transfecting a cell line with a gene that specifies the activity. Furthermore, it is possible to generate a stable transfected cell line that expresses the wild-type F protein in the virus envelope, thereby providing infectious particles (themselves not provided

with genomic information encoding wild-type F protein) with means to enter a cell. Rescue of infectious lentogenic virus is also possible by infection of transfected cells with an NDV helper virus. An essential requirement for such a helper virus would be that it can be selected against, for instance by means of neutralizing antibodies which eliminate the helper virus but which do not react with the lentogenic virus.

**[0083]** Finally, one may construct a stably transfected cell line that expresses one, two, or all of the three essential NDV proteins, NP, P, and L. Such cell lines require the co-expression of only a subset of the three essential proteins or no co-expression at all for supporting generating infectious copy virus.

**[0084]** In a preferred embodiment, the invention provides a method wherein the cells used for transfecting are derived of chicken primary or secondary cells or cell lines. The description provides for example CER or CEF cells, which, as most *in vitro* cells in general, lack the appropriate proteases that are required to cleave the Fo protein of NDV, for example of strain LaSota. However, cells derived from for example other birds can also be used.

**[0085]** The invention further provides a method to generate infectious copy NDV comprising transfecting cells with cloned full-length or genomic-length cDNA of the virus as for example identified in SEQ ID NO:134 and further comprising incubating the cells in growth medium comprising proteolytic activity allowing cleavage of the Fo protein of the virus, further comprising recovering infectious virus by culturing the cells and inoculating material derived from the cultured cells into the allantoic cavity of embryonated eggs. The material for example comprises (harvested or freeze-thawed) cells or cell debris or supernatant derived from the cell culture.

**[0086]** For example, the description describes a method to recover infectious virus, wherein the supernatant of transfected CEF monolayers was inoculated into the allantoic cavity of embryonated eggs. Four days later the allantoic fluid was harvested, analyzed in a hemagglutination assay, and passaged further in eggs.

**[0087]** In addition, the invention provides a method further comprising passaging the infectious copy NDV by harvesting allantoic fluid and re-inoculating embryonated eggs.



**[0088]** In a preferred embodiment, a method is provided wherein the virus is a lentogenic virus, for example derived from an avirulent field-case of NDV or from a vaccine strain of NDV, such as the LaSota strain of NDV. Furthermore, a method is provided to modify an avian-paramyxovirus genome by means of genetic modification which allows the introduction of one or more mutations, deletions, and/or insertions or other modifications. For example, method is provided to attenuate or modify the virulence of avian paramyxovirus by modifying cDNA, for example encoding a viral protein, such as the V protein, and cloning the modified cDNA into full-length cDNA and generating infectious copy virus from the full-length cDNA, thereby generating new NDV strains or new attenuated live vaccines with improved properties.

**[0089]** Apart from attenuation by modification of gene products it is also possible to attenuate avian paramyxovirus by modification of nucleotide sequences which are involved in transcription and/or replication. Such modifications result in attenuated strains which express wild-type-like F proteins which are cleavable both *in vitro* and *in vivo* in a wide range of cells and as a result are more immunogenic than the classical vaccine strains.

**[0090]** In a preferred embodiment, the invention provides a method to attenuate or modify the virulence of an avian paramyxovirus such as a NDV, comprising modifying a protease cleavage site of a viral protein by modifying cDNA encoding the cleavage site, further comprising cloning the cDNA into genomic length cDNA of, for example, NDV and generating infectious copy NDV. The cleavage site is for example a protease cleavage site in the F or HN protein of NDV. Attenuation is in general restricted to reduction of virulence, however, it is now also possible to use a relatively a-virulent strain of NDV and provide the progeny of such a strain with increased virulence, for example by providing it with an increased tendency to replicate in a specified cell-type. It is now thus possible to assign distinct virulence attributes to NDV.

**[0091]** The invention provides a method to antigenically modify avian paramyxovirus such as a NDV, comprising modifying cDNA encoding at least a part of a viral protein harboring at least one immunodominant epitope, further comprising cloning the cDNA into genomic length cDNA of NDV and generating infectious copy NDV.

**[0092]** For example, the invention provides a method to (further) modify NDV, for example using a method to produce an infectious copy of NDV (vaccine) which has been

provided, a method to produce a recombinant marker NDV vaccine is provided, a marker vaccine that contains the fullest possible or needed immunological spectrum of antigenically relevant NDV epitopes, and yet is serologically distinct from wild-type NDV because a distinct, serologically relevant epitope or marker has been removed by recombinant techniques. The invention provides a method to modify the antigenic make-up of avian paramyxovirus such as NDV, thus allowing the generation of, for example, a live NDV marker vaccine which can be serologically distinguished from avian-paramyxovirus field strains.

**[0093]** In one embodiment, the invention provides infectious copy NDV wherein the HN protein of NDV has been modified by recombining cDNA encoding a part of the HN protein with cDNA encoding a part of HN protein derived from an avian paramyxovirus, for example, type 2 or type 4. The hybrid HN protein serves as a serological marker for the infectious copy NDV strain thus obtained or can serve to change the tropism of the avian paramyxovirus to other cells and/or tissues. These, so called “marker strains” as provided by the invention allow the generation of vaccines which are an invaluable tool to assess the prevalence of NDV in commercial flocks around the world. Furthermore, the large-scale application of such marker vaccines will lead to the complete eradication of NDV by a process of intensive screening and stamping out of infected flocks.

**[0094]** Furthermore, a method is provided to generate an infectious copy NDV strain which expresses one or more antigens from other pathogens and which can be used to vaccinate against multiple diseases. Such an infectious copy NDV virus for example comprises a heterologous cDNA encoding a heterologous protein obtained from for example Avian Influenza (AI) (Hemagglutinin (H5 and H7) and Neuraminidase), Avian leukosis virus (ALV) (env protein (gp85)), Chicken anemia virus (CAV) (VP1+VP2), Marek’s disease virus (MDV) (glycoprotein B (gB), gH), Infectious laryngotracheitis virus (ILT) (gB, gH, gD), Infectious bursal disease virus (IBDV) (VP2 and VP3), Turkey rhinotracheitis virus (TRT) (fusion (F) protein), Avian paramyxovirus-2, -3, -6 (PMV) (F-protein, Hemagglutinin neuraminidase (HN), or others, Infectious bronchitis virus (IBV) (peplomer protein, nucleoprotein), Reoviruses (sigma protein), Adenoviruses, Pneumoviruses, *Salmonella enteritidis*, *Campylobacter jejuni*, *Escherichia coli*, *Bordetella avium* (formerly *Alcaligenes faecalis*), *Haemophilus paragallinarum*, *Pasteurella*

*multocida*, *Ornithobacterium rhinotracheale*, *Riemerella* (formerly *Pasteurella*) *anatipestifer*, *Mycoplasma* (*M. gallisepticum*, *M. synoviae*, *M. mereagris*, *M. iowae*), or *Aspergilli* (*A. flavus*, *A. fumigatus*).

**[0095]** The invention herewith provides avian paramyxovirus or strains derived thereof which can be used as a vaccine vector for the expression of antigens from other poultry pathogens. Several properties make NDV an ideal vaccine vector for vaccination against respiratory or intestinal diseases. 1) NDV can be easily cultured to very high titers in embryonated eggs. 2) Mass culture of NDV in embryonated eggs is relatively cheap. 3) NDV vaccines are relatively stable and can be simply administered by mass application methods such as by drinking water or by spraying or aerosol formation. 4) The natural route of infection of NDV is by the respiratory and/or intestinal tracts that are also the major natural routes of infection of many other poultry pathogens. 5) NDV can induce local immunity despite the presence of circulating maternal antibody.

**[0096]** It has been shown that NDV has potent antineoplastic, as well as immune-stimulating properties (for a review see Schirmacher et al., 1998) [Schirmacher, V., Ahlert, T., Steiner, H.-H., Herold-Mende, C., Gerhards, R. and Hagmüller E. (1998) Immunization with virus-modified tumor cells. *Seminars in Oncology* 25: 677-696]. Although NDV does not seem to be able to replicate productively in normal human cells, a selective NDV-mediated killing of human cancer cells was noted. After local NDV therapy, viral oncolysis and complete remissions of human tumor xenografts were observed in nude mice. This has led to the use of NDV for tumor therapy. However, a problem is that such application may be restricted to local treatment.

**[0097]** NDV infection induces interferons, chemokines, and other potentially important gene products, and introduces pleiotropic immune-stimulatory properties into tumor cells. This concept has been used for the production of autologous tumor cell vaccines consisting of fresh operative specimens that have been infected with NDV. This type of vaccine is called autologous tumor vaccine-NDV or ATV-NDV (Schirmacher et al., 1998). The NDV-infected cells are inactivated by gamma-irradiation which prevents cell division but which still allows replication of NDV in the cytoplasm of infected cells. After inoculation of patients with

ATV-NDV, T-cells are recruited through NDV-induced chemokines. Some of these T-cells may express a T-cell receptor that can interact with peptides from tumor-associated antigens in complex with major histocompatibility complex class I molecules at the cell surface. This interaction results in the induction of a cytotoxic T-cell response which results in specific killing of autologous tumor cells.

**[0098]** The invention provides that the repertoire and amount of chemokines and immune-stimulatory proteins induced by NDV infection are modulated. The present invention provides a method for generating recombinant NDV that has been modified to incorporate and express (a) heterologous gene(s). Such recombinant NDV may be used to modify the repertoire and amount of immune-stimulatory proteins in infected cells. In one embodiment, the invention provides a recombinant NDV that incorporates and expresses genes encoding human interferons, chemokines or other immune-stimulatory proteins. The recombinant NDV is used for the production of ATV-NDV which is more potent than conventional ATV-NDV. For example: cytokines IFN- $\alpha$ , - $\beta$ , TNF- $\alpha$ , IL-1, IL-6; chemokines RANTES, IP-10; other genes such as HSP, ACTH, endorphin, iNOS, EPA/TIMP, NF $\kappa$ B.) The pleiotropic immune-stimulatory properties of NDV may also be used as an adjuvant for vaccination of animals and humans against infectious diseases. In one embodiment of the invention, foreign genes encoding (a) relevant antigen(s) of (an) infectious agent(s) are introduced in the NDV genome and the simultaneous expression of the antigen(s) and the immune-stimulatory proteins by infected cells may induce a potent immune response against the infectious agent. In another embodiment of the invention, the immune-stimulating properties of NDV may be further enhanced by using NDV recombinants that simultaneously express antigens and specific immune-stimulatory proteins. In a preferred embodiment, the invention is used to generate an AIDS (acquired immune-deficiency syndrome) vaccine by using NDV recombinants that express relevant antigens of human immune-deficiency virus HIV), either alone or in combination with immune-stimulatory proteins.

**[0099]** NDV are also used as an adjuvant for vaccination of animals and humans against infectious diseases. In one embodiment of the invention, heterologous or foreign genes encoding (a) relevant antigen(s) of (an) infectious agent(s) are introduced in the NDV genome

and the simultaneous expression of the antigen(s) and the immune-stimulatory proteins by infected cells may induce a potent immune response against the infectious agent. In another embodiment of the invention, the immune-stimulating properties of NDV are further enhanced by using NDV recombinants that simultaneously express antigens and specific immune-stimulatory proteins. In a preferred embodiment, the invention is used to generate an AIDS (acquired immune-deficiency syndrome) vaccine by using NDV recombinants that express relevant antigens of human immune-deficiency virus (HIV), either alone or in combination with immune-stimulatory proteins.

**[00100]** Also, a method is provided to generate a conditional lethal NDV deletion mutant which can be used as self-restricted non-transmissible (carrier) vaccine. An NDV deletion mutant was generated which is unable to express the matrix (M) protein which is involved in budding of NDV at the inner cell membrane. The invention provides for example a phenotypically complemented NDV strain that is unable to express the M protein which is still able to infect cells and spread by means of cell-to-cell transmission. However, the mutant virus is unable to generate infectious progeny on non-complementing cells. This shows that phenotypically complemented NDV deletion mutants can be used as safe self-restricted vaccines which are unable to spread into the environment. Such a non-transmissible vaccine combines the most important advantage of live vaccines, *i.e.*, efficacy, with the most important advantage of killed vaccines, *i.e.*, safety.

**[00101]** The invention provides NDV, or strains derived thereof, for example by passaging or further cultivation in embryonated eggs or appropriate cells, that is derived from infectious copy virus obtainable by a method provided by the invention.

**[00102]** For example, NDV is provided that has been modified in at least one way to generate infectious copy NDV which is attenuated, modified in virulence, antigenically modified, expressing a heterologous antigen or are non-transmissible, or combinations thereof.

**[00103]** Herewith the invention provides NDV vaccines, characterized for example by carrying distinct virulence attributes or distinct antigenic characteristics, be it for marker vaccine purposes and/or for expressing heterologous antigens derived from other pathogens, be it in transmissible and/or non-transmissible form.

**[00104]** Such a vaccine can be a killed or a live vaccine, preferably, such a vaccine is a live vaccine, however, killed vaccines as provided by the invention are beneficial under those circumstances where a live vaccine is not or only little applicable, for example because of trade restrictions or other conditions set by disease controlling authorities.

**[00105]** The invention herewith also provides a diagnostic method, and corresponding test kit, to detect antibodies against the serologically relevant immunodominant epitope or marker, therewith providing methods and means to execute a method for control and/or eradication of NDV and/or other poultry diseases in poultry. The invention provides new and effective vaccines which can be serologically discriminated from field-viruses and old-type vaccines. Such new vaccines, called NDV marker vaccines, provide the fullest possible immunological spectrum of antigenically relevant NDV epitopes, and yet are serologically distinct from wild-type NDV by applying accompanying diagnostic methods and kits.

**[00106]** The invention provides a method for distinguishing unvaccinated animals or animals vaccinated with a NDV vaccine according to the invention from animals infected with wild-type NDV or vaccinated with an unmodified mesogenic or lentogenic NDV-vaccine strain comprising taking a least one sample (such as serum, blood, eggs or eye fluid) from the animal and determining in the sample the presence of antibodies directed against an immunodominant epitope or marker expressed by the wild-type or unmodified NDV but not by a vaccine according to the invention.

**[00107]** The invention provides a method wherein the antibodies are directed against the HN or F protein of NDV, for example a hybrid protein as described in the experimental part as this description. The invention provides for example a diagnostic method wherein the animal is selected from the group composed of poultry, preferably of chickens.

**[00108]** The invention also provides a diagnostic kit for use in a method to serologically distinguish between animals. In one embodiment of the invention, a simple and rapid hemagglutination-inhibition (HI) test is used to distinguish between vaccinated animals and infected animals. Animals vaccinated with a marker vaccine in which the complete globular head of HN of NDV has been replaced with the corresponding part of HN of another serotype

will not induce antibodies to HN of NDV and therefore will not inhibit hemagglutination of erythrocytes by NDV virions.

[00109] By using marker vaccine virions in the HI test, antibodies against the hybrid HN protein is detected and may be used as a measure for the efficacy of vaccination. As an alternative, an ELISA that detects antibodies against the F protein of NDV is used to measure the efficacy of vaccination.

[00110] Apart from the HI test, an ELISA can be used to determine the presence of antibodies against HN of NDV. The antigen to be used in such a test is for example HN of NDV that is expressed by recombinant DNA techniques or a conserved peptide from HN of NDV.

[00111] A blocking ELISA may also be used. In this case one or more monoclonal antibodies against conserved epitopes of HN of NDV are used to determine whether competing antibodies are present in samples from vaccinated animals. The ELISA tests can advantageously be used if the marker vaccine contains a chimeric HN protein only or when a few epitopes of HN of NDV are replaced.

[00112] The invention is further explained in the experimental part of this description without limiting the invention thereto.

#### DESCRIPTION OF THE FIGURES

[00113] FIG. 1. Transcription vector pOLTV5 is a derivative of the transcription vector described by Pattnaik et al. (1992). See text for details of the construction. The plasmid contains the T7 DNA-dependent RNA polymerase promoter (shown in boldface) followed by unique *Stu*I and *Sma*I restriction sites and the autocatalytic ribozyme from hepatitis delta virus (HDV). DNA fragments can be cloned between the *Stu*I and *Sma*I sites and can be transcribed either *in vitro* or *in vivo* by using T7 RNA polymerase. The 5' end of the resulting transcripts contains two extra G-residues which are not encoded by the insert. Due to the action of the ribozyme, the 3' end of the transcripts exactly corresponds to the last nucleotide of the insert.

[00114] FIG. 2. Structure of the minigenome plasmids pOLTV535 (Part A of FIG. 2) and pOLTV553 (Part B of FIG. 2). The minigenome plasmids are based on transcription plasmid pOLTV5 (cf. FIG. 1) and contain the 3'-region (nt 1-119) and 5'-region (nt

14970-15186) of NDV strain LaSota flanking the gene encoding secreted alkaline phosphatase (SEAP). Transcription of pOLTV535 by T7 RNA polymerase yields antigenomic RNA (or ([+]-RNA) whereas transcription of pOLTV553 yields genomic RNA (or [-]-RNA). The start (S) and end (E) boxes, which are viral transcription initiation and termination signals, are indicated. The start codon of the SEAP gene is underlined. The sequences of the insertions (N0-N5) in the ClaI site which generate minigenome plasmids which each differ 1 nt in length (pOLTV535N0-N5 and pOLTV553N0-N5, respectively) are also shown.

**[00115]** FIG. 3. FIG. 3A depicts the overall strategy used for the assembly of the full-length NDV cDNA from subgenomic overlapping cDNA clones. The cDNA was assembled in plasmid pOLTV535 which already contained the 3' and 5' ends of NDV strain LaSota (cf. FIG. 2). The resulting plasmid, designated pNDFL+, was used for the generation of infectious NDV.

**[00116]** FIG. 3B depicts the detailed cloning procedure for the assembly of the full-length NDV cDNA from subgenomic overlapping cDNA clones. Cm denotes the chloramphenicol-resistance gene which was temporarily introduced as a phenotypical tag (see text for details).

**[00117]** FIG. 3C depicts the detailed cloning procedure for the generation of genetically modified full-length NDV cDNA. The modification consists of 3 nucleotide changes which were introduced in the F gene and which result in the modification of the amino acid sequence of the proteolytic cleavage site of the F protein (see text for details).

**[00118]** FIG. 4A depicts the pOLTV535-series. Transcription by means of T7 RNA polymerase yields antigenomic RNA (or [+]-RNA) which can be directly translated into SEAP protein by the cell. After infection of cells by helper virus (or after co-transfection of plasmids encoding NP, P, and L), the antigenomic RNA is used by the viral polymerase complex for the synthesis of genomic RNA (or [-]-RNA). The genomic RNA is then used by the viral polymerase complex for the synthesis of both mRNA (by using the specific transcription start [S] and end [E] boxes) and antigenomic RNA.

**[00119]** FIG. 4B depicts the pOLTV553-series. Transcription by means of T7 RNA polymerase yields genomic RNA (or [-]-RNA) which cannot be translated into SEAP protein.



After infection of cells by helper virus (or after co-transfection of plasmids encoding NP, P and L), the genomic RNA is used by the viral polymerase complex for the synthesis of both mRNA (by using the specific transcription start [S] and end [E] boxes) and antigenomic RNA.

**[00120]** FIG. 5. Alignment of nucleic acid sequences 5'-terminal ends of NDV LaSota and other paramyxoviruses are given as sequence comparison of NDV across the four members of the *Rubulavirus* genus, three members of the *Paramyxovirus* genus, and three members of the *Morbillivirus* genus. The sequences are presented from the L gene end box to the 5' end (3'. 5' cDNA). NDV, NDV (SEQ ID NO:143); hPIV2, human parainfluenza virus 2 (SEQ ID NO:144); MuV, mumps virus (SEQ ID NO:145); SV5 (SEQ ID NO:147) and SV41 (SEQ ID NO:146), simian virus 5 and 41, respectively; SeV, sendai virus (SEQ ID NO:140); bPIV3 (SEQ ID NO:140) and hPIV3 (SEQ ID NO:141) bovine and human parainfluenza virus, respectively; CDV, canine distemper virus (SEQ ID NO:137); MeV measles virus (SEQ ID NO:138); RPV, rinderpest virus (SEQ ID NO:139). Nucleotide (nt) sequences of the entire genomes were obtained as follows (accession no.): NDV (AF077761); hPIV2 (X57559); MuV (AB000388); SV5 (AF052755); SV41 (X64275); bPIV3 (D84095); hPIV3 (Z11575); CDV (L13194); MeV (X16565); RPV (Z30697).

## DETAILED DESCRIPTION OF THE INVENTION

### EXPERIMENTAL PART

#### MATERIALS AND METHODS

**[00121]** Standard cloning procedures were carried out according to Sambrook et al. (1989) unless stated otherwise. All constructions involving DNA fragments which were generated by means of the polymerase chain reaction (PCR) were verified by sequence analysis. In the primer sequences given below, the underlined nucleotides correspond to NDV sequences and the position within the NDV genome is indicated. The nucleotide sequence of restriction sites which were used for cloning are indicated in boldface.

## Cells and viruses

**[00122]** CER cells (Smith et al., 1976) were grown in GMEM/EMEM (1:1) containing 5% fetal calf serum and 2% of an antibiotic mix that contained 1000 U/ml Penicillin, 1000 µg/ml Streptomycin, 20 µg/ml Fungizone, 500 µg/ml Polymixin B, and 10 mg/ml Kanamycin. QT35 cells (Moscovici et al., 1977; Cho, 1982) were grown in medium supplied by GibcoBRL/Life Technologies (cat. no. 041-91536; proprietary composition of Fort Dodge Labs, Ft. Dodge, Iowa) supplemented with 5% FCS and 2% antibiotic mix. QM5 cells (Antin and Ordahl, 1991) were grown in M199 medium supplemented with 10% tryptose phosphate broth, 10% FCS and 2% antibiotic mix.

**[00123]** NDV strain LaSota was obtained from ATCC (ATCC VR-699) and was passaged two times in embryonated eggs. Before we started with the construction and cloning of cDNA, the virus was plaque purified by three rounds of plaque purification on primary chicken embryo fibroblasts (CEF). To this end, the virus was titrated on CEF cells cultured in GMEM/EMEM (1:1) containing 5% fetal calf serum, 2% antibiotic mix, 5% allantoic fluid, 30 mM MgCl<sub>2</sub>, 200 µg/ml DEAE-dextran (Sigma) and 0.8% agar Nobel (Difco). Virus from the third round of plaque purification (designated clone E13-1) was grown in embryonated eggs and four days after inoculation the allantoic fluid was harvested and stored in aliquots at -70°C. The fowl pox recombinant virus fpEFLT7pol (Britton et al., 1996; hereafter called FPV-T7), which expresses T7 RNA polymerase, was a kind gift of Dr. Michael Skinner and was grown on QT35 cells.

## Isolation of viral RNA

**[00124]** All manipulations were carried out in RNase-free glassware or plastics and all solutions were made up with RNase-free water which was treated with 1% diethyl-pyrocabonate (DEPC) and sterilized by autoclaving. Virus was pelleted from allantoic fluid by centrifugation at 21,000 rpm for 70 minutes in a Beckman SW40 rotor at 4°C. The pellet was resuspended in homogenization buffer (50 mM Tris-HCl pH 7.5, 50 mM NaCl, 5 mM EDTA, 0.5% SDS) and treated with Proteinase K (200 µg/ml) for 90 minutes at 37°C during constant agitation. The lysate was extracted two times with an equal volume of

phenol/chloroform (1:1) pH 5.4 and once with an equal volume of chloroform. The viral RNA was precipitated from the aqueous phase by the addition of 0.1 volume of 3M NaOAc pH 5.3 and 2.5 volumes of 100% ethanol. The precipitate was collected by centrifugation, washed once with 70% ethanol, resuspended in water, and stored in aliquots at -70°C.

#### Reverse transcription

**[00125]** Viral RNA (1.5 µg) was mixed with 500 ng of primer in a volume of 12 µl and incubated for 10 minutes at 70°C. Four µl of 5x RT buffer (250 mM Tris-HCl, pH 8.3, 375 mM KCl, 15 mM MgCl<sub>2</sub>; GibcoBRL/Life Technologies), 2 µl 0.1 M DTT and 2 µl 10 mM dNTP's (2.5 mM each) was added and the mixture was incubated for 2 minutes at 42°C. Reverse transcription was performed in a final volume of 20 µl by the addition of 200 Units of reverse transcriptase (Superscript II; GibcoBRL/Life Technologies) followed by incubation for 60 minutes at 42°C.

#### Polymerase Chain Reaction (PCR)

**[00126]** All PCR reactions which were used to determine the 3' and 5' end of the NDV genome (see below) were carried out by using Taq DNA Polymerase (Perkin Elmer). For the cloning of individual NDV genes or large subgenomic cDNA's, either the proofreading DNA polymerase Pwo, or mixtures of Taq and Pwo (Expand High Fidelity Kit or Expand Long Template Kit) were used according to the instructions of the supplier (Boehringer Mannheim). All samples were incubated for 2 minutes at 94°C before the start of the indicated number of PCR cycles. After the indicated number of PCR cycles, the samples were incubated at the elongation temperature for at least 3x the duration of the elongation time of the PCR cycle. PCR fragments were purified directly by using the High Pure PCR Product purification Kit (Boehringer Mannheim) or after agarose gel electrophoresis by using the QiaexII extraction kit (Qiagen) essentially as described by the suppliers.

## Sequence analysis

[00127] All sequences were determined by using the PRISM Ready Reaction Dye Deoxy Terminator Cycle Sequencing Kit (Perkin Elmer). Reaction mixtures (5 µl) were subjected to 25 cycles of linear amplification (10 seconds at 94°C, 5 seconds at 50°C, and 4 minutes at 60°C) in a GeneAmp2400 thermocycler. Subsequently, the reaction mixtures were precipitated with ethanol, washed once with 70% ethanol, resuspended in 15 µl TSR buffer (Perkin Elmer) and heated for 2 minutes at 94°C before being loaded on an Applied Biosystems AB310 automatic sequencer.

[00128] The nucleotide sequences of the primers which were used to sequence the complete genome of NDV strain LaSota were either derived from published sequences or from sequences established during this sequencing project. The primers are shown in Table 1.

## Cloning and sequencing of the 3' and 5' termini of the genome of NDV strain LaSota

[00129] The nucleotide sequence of the 3' and 5' termini of the NDV genome were determined by using RACE procedures (rapid amplification of cDNA ends). NDV RNA was used in a reverse transcription reaction in a final volume of 20 µl by using primer p360 (5'-GGCGATGTAATCAGCCTAGTGCTT-3' (SEQ ID NO:47); nt 14756-14779) which was derived from the published sequence of the L gene of NDV (Yusoff et al., 1987). The single-stranded cDNA (2.5 µl of the RT mixture) was added to 8 pmol anchor primer ALG3 (5'-CACGAATTC<sup>ACTATCGATTCTGGATCCTTC</sup>-3' (SEQ ID NO:83)) and ligated overnight at room temperature in 20 µl of a reaction mixture containing 50 mM Tris-HCl, pH 8.0, 10 mM MgCl<sub>2</sub>, 10 µl/ml BSA, 25% PEG, 1 mM HCC, 20 uM ATP and 10 units of T4 RNA ligase (New England Biolabs) as described by Tessier et al. (1986). One µl of the ligation reaction was used as template in a PCR reaction by using primers p375 (5'-CAATGAATTCAAAGGA TATTACAGTAACT-3' (SEQ ID NO:84); nt 14964-14983) and ALG4 (5'-GAAGGATCCAGAATCGATAG-3' (SEQ ID NO:85)). The latter primer is complementary to anchor primer ALG3 (SEQ ID NO:83). The PCR conditions (40 cycles) were as follows: 1 minute at 94°C, 1 minute at 55°C, and 2 minutes at 72°C. The PCR products were purified and cloned in T-vector pBluescriptII-TSK (Ichihara and Kurosawa, 1993). Alternatively, the purified

PCR products were treated with Klenow DNA polymerase I to create blunt ends and cloned in the HindII site of plasmid pGEM4Z (Promega). Thirteen independent clones (8x pBluescriptII-TSK and 5x pGEM4Z) were sequenced to determine the nucleotide sequence of the 5' end of the genome of NDV strain LaSota. The nucleotide sequence of the 3' end was determined by two independent methods. In method I, primer ALG3 (SEQ ID NO:83) was ligated to the 3' end of the viral RNA by using T4 RNA ligase as described by Schütze et al. (1995). The reaction mixture (final volume 10 µl) contained 2.5 µg NDV RNA, 100 pmol ALG3 (SEQ ID NO:83), 1 µl 10x T4 RNA ligase buffer (500 mM Tris-HCl, pH 7.8, 100 mM MgCl<sub>2</sub>, 100 mM DTT, 10 mM ATP), 1 µl DMSO, 1 µl 10 uM hexamine-cobalt chloride, 1 µl RNasin (Promega) and 10 units of T4 RNA ligase (New England Biolabs). The mixture was incubated overnight at room temperature and 5 µl of the ligation reaction was used as template in a reverse transcription reaction by using ALG4 (SEQ ID NO:85) as primer. One µl of the RT-reaction was used in a PCR reaction by using primers ALG4 (SEQ ID NO:85) and p376 (5'-GAGCCTTAAGGAGCTGCTCGTACTGATC-3' (SEQ ID NO:86); nt 137-164) which was derived from the published sequence of the 3' end of NDV (Ishida et al., 1986). The PCR conditions were as described above for the 5'-RACE. In method II, the 3' and 5' ends of the viral NDV RNA were ligated to each other by using T4 RNA ligase using the same conditions as described above for method I. Five µl of the ligation mixture was used as template in a reverse transcription reaction by using primer p360 (SEQ ID NO:47). One µl of the RT-reaction was used in a PCR reaction by using primers p375 (SEQ ID NO:84) and p376 (SEQ ID NO:86) and PCR conditions described above for the 5'-RACE. The PCR products were treated with Klenow DNA polymerase I to create blunt ends and cloned in the HindII site of plasmid pGEM4Z (Promega). Ten independent clones (4 from method I and 6 from method II) were sequenced to determine the nucleotide sequence of the 3' end of the genome of NDV strain LaSota.

#### Construction of transcription vector

**[00130]** A low-copy-number transcription vector was constructed by using plasmid pOK12 (Vieira and Messing, 1991) as the basic replicon. Plasmid pOK12 was digested with PvuII and the DNA fragment containing the replication origin and the Kanamycin-resistance

gene was isolated. This DNA fragment was ligated to an Eco47III-AflIII fragment (the AflIII site was made blunt by using Klenow DNA polymerase I) from transcription vector 2.0 (a generous gift of Dr. Andrew Ball; Pattnaik et al., 1992). From the resulting plasmid an XbaI-NheI fragment was deleted to eliminate as much unique restriction sites as possible. The resulting plasmid was designated pOLTV5 (FIG. 1). Transcription vector pOLTV5 contains the T7 DNA-dependent RNA polymerase promoter followed by unique StuI and SmaI restriction sites, the autocatalytic ribozyme from hepatitis delta virus (HDV) and the transcription termination signal from bacteriophage T7. DNA fragments cloned between the StuI and SmaI restriction sites can be transcribed either *in vitro* or *in vivo* by using T7 RNA polymerase. After transcription, the 5' end of the resulting transcripts contains two G residues encoded by the plasmid. Due to the autocatalytic action of the HDV ribozyme, the 3' end of the transcripts corresponds to the exact terminal nucleotide of the cloned DNA fragment (Pattnaik et al., 1992).

#### Construction of minigenome plasmids

**[00131]** In order to examine the requirements for replication and transcription of NDV, minigenome plasmids were constructed which contained the 3'- and 5'-terminal regions of NDV flanking a reporter gene that replaced all NDV genes (FIG. 2). DNA fragments corresponding to the 3'- and 5'-terminal regions of NDV were generated by means of PCR by using Pwo DNA polymerase (30 cycles; 15 seconds at 94°C, 30 seconds at 50°C, and 30 seconds at 72°C) and using plasmids containing the 3'- and 5'-RACE fragments as templates (see above).

**[00132]** The 3'-region (nt 1-119) was generated by using primers 3UIT (5'-ACCAAACAGAGAATCCGTGAGTTACGA-3' (SEQ ID NO:1); nt 1-27) and SEAP3 (5'-ATCGATACTGGTCAGCATGCTGGCAGAAGGCTTTCTCG-3' (SEQ ID NO:87); nt 102-119). The 5'-region (nt 14973-15186) was generated by using primers SEAP5 (5'-GCATGCTGACCAGTATCGATATTACAGTAACTGTGACT-3' (SEQ ID NO:88); nt 14973-14990) and 5NDV (5'-ACCAAACAAAGATTTGGTGAATGACGA-3' (SEQ ID NO:79); nt 15158-15186). The two DNA fragments were joined in an overlap PCR (the overlap is shown in italics in the primer sequences shown above) by using primers 3UIT (SEQ ID NO:1) and 5NDV (SEQ ID NO:79). The resulting DNA fragment, which is a fusion of the 3' and 5'

ends of NDV separated by 20 nucleotides, was phosphorylated by treatment with T4 polynucleotide kinase and cloned in both orientations in transcription plasmid pOLTV5 (FIG. 1) which was cleaved with *Stu*I and *Sma*I and dephosphorylated with calf intestinal phosphatase (Boehringer Mannheim). Finally, the SEAP gene (encoding secreted alkaline phosphatase) was recovered from plasmid pSEAP-Basic (Clontech) by digestion with *Sph*I and *Cla*I, and cloned between the *Sph*I and *Cla*I sites between the 3' and 5' ends of NDV. The resulting plasmids were designated pOLTV535 and pOLTV553, respectively. *In vivo* or *in vitro* transcription using T7 RNA polymerase of plasmid pOLTV535 gives rise to antigenomic RNA ([+]-RNA), whereas transcription of plasmid pOLTV553 gives rise to genomic RNA ([-]-RNA).

**[00133]** Plasmids pOLTV535N0 to -N5 and pOLTV553N0 to -N5 were generated by inserting self-complementary oligonucleotides in the *Cla*I site located between the SEAP gene and the 5' end of NDV in pOLTV535 and pOLTV553, respectively (see FIG. 2). The oligonucleotides used were: N0, 5'-CGCGAGCTCG-3' (SEQ ID NO:89); N1, 5'-CGCGAGSCTCG-3' (SEQ ID NO:90); N2, 5'-CGCGAGCGCTCG-3' (SEQ ID NO:91); N3, 5'-CGCGAGCWGCTCG-3' (SEQ ID NO:92); N4, 5'-CGCGAGCATGCTCG-3' (SEQ ID NO:93); N5, 5'-CGCGAGCASTGCTCG-3' (SEQ ID NO:94) (W = A or T; S = C or G).

#### Modification of the T7 promoter in plasmids pOLTV535 and pOLTV553

**[00134]** To generate *in vitro* or *in vivo* transcripts containing the authentic 5'- and 3'-terminal ends of NDV, the T7 promoter in plasmids pOLTV535 and pOLTV553 was modified such that transcription would start at the first nucleotide of the 3'- or 5'-terminal ends of NDV.

**[00135]** Primers were designed which contained, 1) a *Bgl*II-restriction site, 2) the sequence of the T7 promoter (shown in italics) which was modified such that the two G residues at the end of the T7 promoter were replaced by an A residue, and 3) the 3' (nt 1-21) or 5' end (nt 15164-15186) of NDV. Primers BGL3F2 (5'-GATATGGCCATTCAGGCTTAATACGACT CACTATAACCAAACAGAGAATCCGTGAG-3' (SEQ ID NO:95)) and SEAP3 (SEQ ID NO:87) (see above) were used to generate a DNA fragment containing the modified T7 promoter and the entire 3' end of NDV up to the start of the SEAP gene in pOLTV535. Similarly, a DNA fragment was generated containing the modified T7 promoter and the entire 5' end of NDV up to

the end of the SEAP gene in pOLTV553 by using primers BGL5F2 (5'-GATATGGCCATTCAGGCTTAATACGACTCACTATAACCAAACAAAGATTTGGTGAATG-3' (SEQ ID NO:95)) and SEAP5 (SEQ ID NO:88). The resulting fragments were digested with BglII and SphI (3' end) or BglII and ClaI (5' end), respectively, and used to replace the BglII-SphI fragment in pOLTV535, or the BglII-ClaI fragment in pOLTV553. The resulting plasmids were designated pOLTV735 and pOLTV753, respectively. Plasmids pOLTV735N3 and pOLTV753N3 were generated by inserting a self-complementary oligonucleotide (5'-CGCGAGCWGCTCG-3' (SEQ ID NO:92); W = A or T) in the ClaI site located between the SEAP gene and the 5' end of NDV in pOLTV735 and pOLTV753, respectively.

#### Construction of SEAP-reporter plasmids

**[00136]** Plasmid pCIneoSEAP was constructed by cloning an XhoI-ClaI fragment (ClaI site was made blunt by using Klenow DNA polymerase I) containing the SEAP gene from plasmid pSEAP-Basic (Clontech) between the XhoI and SmaI sites of the eukaryotic-expression vector pCIneo (Promega). The latter plasmid contains the human cytomegalovirus (hCMV) promoter in addition to the bacteriophage T7 promoter. In order to examine and quantify SEAP expression by transcripts generated from the T7 promoter only, another plasmid was constructed which lacked the hCMV promoter. To this end, the hCMV promoter was deleted from pCIneo by partial digestion with HindIII followed by complete digestion with BglII. The DNA fragment (nt 756-5469 according to numbering of Clontech) from which the hCMV promoter was deleted was isolated, treated with T4 DNA polymerase to generate blunt ends and recircularized by using T4 DNA ligase. The resulting plasmid was designated pCIneoD. Finally, the SEAP gene was recovered from pSEAP-Basis as a MluI-AccI fragment and cloned in pCIneoD between the MluI and ClaI sites. The resulting plasmid was designated pCIneoD SEAP.

#### Transfections

**[00137]** Cells were seeded in 24-well culture dishes, grown overnight to 60-80% confluency, and infected at an m.o.i. of 1 with FPV-T7 for 1 hour at 37°C. The cells were transfected with 0.5 µg minigenome plasmid DNA by using 3 µl of LipofectAMINE™ and



OptiMem essentially as described by the supplier (GibcoBRL/Life Technologies). After incubation for 4 hours (CER cells) or 16 hours (QM5 cells) at 37°C the cells were either infected with NDV (Dutch virulent isolate no. 152608; 200 µl per well) for 1 hour at a m.o.i. of 5, or left uninfected. The inoculum was aspirated and replaced by 1 ml of complete medium and the cells were further incubated at 37°C.

**[00138]** For co-transfections, cells were grown in 6-well culture dishes and infected with FPV-T7 as described above. The cells were co-transfected with 0.25 µg minigenome plasmid DNA, 0.4 µg pCIneoNP, 0.2 µg pCIneoP and 0.2 µg pCIneoL(c) or pCIneo by using either 8 µl of LipofectAMINE or 9 µl of FuGene™ 6 (Boehringer Mannheim). In order to generate infectious virus, the minigenome plasmid was replaced by a transcription plasmid that contained the full-length NDV cDNA.

#### Quantification of SEAP activity

**[00139]** The amount of SEAP which was secreted into the medium of transfected cells was measured in disposable 96-well plates by using the Phospha-Light™ Chemiluminescent Reporter Assay for Secreted Alkaline Phosphatase kit essentially as described by the supplier (Tropix). Chemiluminescence was quantified by using a liquid scintillation counter (Wallac 1450 microbeta PLUS).

#### Cloning and sequencing of cDNA's spanning the entire genome of NDV strain LaSota

**[00140]** To clone and sequence the entire genome of NDV strain LaSota, large subgenomic cDNA clones were generated by means of RT-PCR and cloned in pGEM-T. First strand cDNA synthesis was performed by using primer 3UIT (SEQ ID NO:1) as described above, and 1 µl of the RTreaction was used in a PCR reaction by using the Expand Long Template PCR kit (Boehringer Mannheim). The PCR consisted of 5 cycles of 10 seconds at 94°C, 30 seconds at 58°C, and 6 minutes at 68°C, followed by 10 cycles of 10 seconds at 94°C, 30 seconds at 58°C, and 6 minutes at 68°C, in which the elongation time at 68°C was increased by 20 seconds per cycle. The PCR fragments were cloned in pGEM-T by using the pGEM-T cloning kit essentially as described by the supplier (Promega). Ligation mixtures were

transformed into *E. coli* strain SURE II (Stratagene). Two independent RT-PCR reactions (A and B) were performed and each yielded a similar set of cDNA clones. The nucleotide sequence of the subgenomic cDNA clones was determined by using NDV-specific primers (Table 1) and by primers flanking the inserts. After comparison of the nucleotide sequence of the A and B series of clones, remaining ambiguities were resolved by sequencing relevant regions of a third independent series of cDNA's (C series). The nucleotide sequence of NDV strain LaSota is shown in SEQ ID NO:134.

**[00141]** Nucleotide sequence (SEQ ID NO: 134) of the genome of NDV strain LaSota and deduced amino acid sequence of the NDV genes. The sequence shown corresponds to the antigenomic strand and is shown in the 5' to 3' direction in the form of ssDNA. The sequence shown in this figure is that of the consensus sequence which was determined by completely sequencing two independent sets of overlapping subgenomic cDNA's which span the entire NDV genome. Remaining ambiguities (probably as a result of PCR errors) were resolved by sequencing relevant regions of a third independent set of clones.

**[00142]** The sequence of the full length cDNA clone pNDFL+ which was assembled from overlapping subgenomic cDNA clones (see FIG. 3), differs from that of the consensus NDV sequence at the following positions (consensus sequence between parentheses): nt 1755, G (A); nt 3766, A (G); nt 5109, G (A); nt 6999, T (C); nt 7056, G (A); nt 9337, G (A); nt 9486, A (T); nt 10195, T (C); nt 13075, A (G). These differences results in 3 amino acid changes (consensus sequence between parentheses): F protein, R<sup>189</sup> (Q); HN protein S<sup>200</sup> (P) L-protein N<sup>369</sup> (I).

#### Construction of a full length genomic cDNA clone of NDV

**[00143]** The full-length NDV cDNA was assembled in transcription plasmid pOLTV5 by using pOLTV535 as the starting plasmid. The DNA fragments were joined at overlaps by using common restriction enzymes as detailed in portion B of FIG. 3. In a series of cloning steps, a plasmid (designated p535-DI) was constructed containing nucleotides 1-3521 and 12355-15186 separated by a ClaI site that was generated by joining the ClaI sites at position 3521 and 12355. In another series of cloning steps, a plasmid (designated pGEM-B) was

constructed which contained part of the NDV genome including nucleotides 3521-12355 (ClaI fragment). To facilitate cloning, the latter ClaI fragment was tagged with the Chloramphenicol-resistance (Cm) gene from plasmid pACYC184 (Chang and Cohen, 1978). To this end, the Cm gene was recovered from pACYC184 by means of PCR by using primers CAT-F (5'-GCGTACGTCTAGACTGGTGTCCCTGTTGATACCGG-3' (SEQ ID NO:96) and CAT-R (5'-GCTCTAGACGTACGACCCTGCCCTGAACCGACG-3' (SEQ ID NO:97). The PCR was carried out with Pwo DNA polymerase and consisted of 30 cycles of 30 seconds at 94°C, 45 seconds at 60°C, and 60 seconds at 72°C. The resulting PCR fragment was digested with BsiWI and cloned in the unique BsiWI site of pGEM-B, yielding PGEM-B (CAT). The ClaI fragment from pGEM-B (CAT) was cloned in the unique ClaI site of p535-DI, yielding pNDFL (CAT). Finally, the Cm gene was removed from this plasmid by digestion with BsiWI followed by religation and transformation of *E. coli* strain DH5a. The resulting plasmid was designated pNDFL+ and contains the entire NDV cDNA sequence cloned between the T7 promoter and the HDV ribozyme in transcription plasmid pOLTV5.

#### Cloning and expression of individual NDV genes

**[00144]** DNA fragments containing each of the NDV LaSota genes were generated by means of RT-PCR and cloned in pCIneo. After cloning, all fragments were sequenced by using primers flanking the inserts and by gene-specific primers. NP gene: Primer 386 (5'-GAGCAATCGAAGTCGTACGGGTAGAAGGTG-3' (SEQ ID NO:98); nt 40-69) was used for reverse transcription. Primers 365 (5'-GTGTGAATTCCGAGTGCGAGCCCCGAAG-3' (SEQ ID NO:99); nt 77-94) and 892 (5'-TTGCATGCCTGCAGGTCAGTACCCCCAGTC-3' (SEQ ID NO:100); nt 1577-1593) were used for PCR by using Pwo DNA polymerase. The following PCR profile (30 cycles) was used; 30 seconds at 95°C, 40 seconds at 65°C, and 45 seconds at 72°C. The resulting DNA fragment was digested with EcoRI and cloned in pCIneo between the EcoRI and SmaI sites. Expression of NP was verified in an immunoperoxidase monolayer assay (IPMA) as described by Peeters et al. (1992) by using monoclonal antibody 38 (Russell et al., 1983). P gene: Primer pRT1 (5'-CAAAGAATTCAGAAAAAAGTACGGGTAGAA-3' (SEQ ID NO:8); nt 1794-1814) was used for reverse transcription. Primers pRT1 (SEQ ID NO:8) and p2

(5'-GCAGTCTAGATTAGCCATTCACTGCAAGGCGC-3' (SEQ ID NO:101); nt 3053-3071) were used for PCR by using Pwo DNA polymerase. The following PCR profile (30 cycles) was used; 30 seconds at 95°C, 40 seconds at 65°C, and 60 seconds at 72°C. The resulting DNA fragment was digested with EcoRI and XbaI and cloned in pCIneo between the EcoRI and XbaI sites. Expression of P was verified in an IPMA by using monoclonal antibody 688 (Russell et al., 1983).

**[00145]** M gene: Primer 3UIT (5'-ACCAAACAGAGAATCCGTGAGTTACGA-3' (SEQ ID NO:1); nt 1-27) was used for reverse transcription. Primers NDV5M (5'-GGGTGCTAGCGGAGTGCCCCAATTGTGCCAA-3' (SEQ ID NO:102); nt 3268-3288) and NDV3M (5'-TCTCCCCGGGGCAGCTTATTTCTTAAAAGGAT-3' (SEQ ID NO:56); nt 4368-4389) were used for PCR by using the Expand High Fidelity kit. The PCR consisted of 10 cycles of 15 seconds at 95°C, 30 seconds at 55°C, and 2 minutes at 68°C, followed by 15 cycles in which the elongation time at 68°C was increased for 20 seconds per cycle. The resulting DNA fragment was treated with T4 DNA polymerase to create blunt ends, digested with NheI, and cloned in pCIneo between the NheI and SmaI sites. Expression of the M protein was verified in an IPMA by using monoclonal antibody 424 (Russell et al., 1983).

**[00146]** F gene: Primer 3UIT (SEQ ID NO:1) (see above) was used for reverse transcription. Primers NDV5F (5'-ACGGGCTAGCGATTCTGGATCCCGGTTGG-3' (SEQ ID NO:15); nt 4508-4526) and NDV3 F (5'-ACTACCCGGGAAACCTTCGTTCTCAT-3' (SEQ ID NO:60); nt 6212-31) were used for PCR by using the Expand High Fidelity kit using the conditions described above for the M gene. The resulting DNA fragment was treated with T4 DNA polymerase to create blunt ends, digested with NheI, and cloned in pCIneo between the NheI and SmaI sites. Expression of the F protein was verified in an IPMA by using monoclonal antibody 8E12A8C3 (ID-DLO, department of Avian Virology).

**[00147]** HN gene: Primer 3UIT (SEQ ID NO:1) was used for reverse transcription. Primers NDV5HN (5'-GTAGGCTAGCAAGAGAGGCCGCCCTCAAT-3' (SEQ ID NO:22); nt 6335-6354) and NDV3HN (5'-CGAGCCCGGGCCGGCATTCGGTTTGATTCTTG-3' (SEQ ID NO:104); nt 8205-8227) were used for PCR by using the Expand High Fidelity kit using the conditions described above for the M gene. The resulting DNA fragment was treated with T4

DNA polymerase to create blunt ends and after digestion with XmaI it was cloned in pCIneo between the blunted (Klenow DNA polymerase) NheI site and the XmaI site. Expression of the HN protein was verified in an IPMA by using monoclonal antibody 86 (Russell et al., 1983).

**[00148]** L gene: The L gene was recovered from cDNA clone pGEM-L7a (Portion A of FIG. 3) by digestion with SacII and SalI. Before digestion with SalI, the SacII site was made blunt by treatment with T4 DNA polymerase. The resulting fragment was cloned in pCIneo between the blunted (Klenow DNA polymerase) NheI site and the SalI site. The 5' untranslated region between the T7 promoter and the ATG start codon of the L gene contained 2 out-of-frame ATG codons which might interfere with correct expression of the L protein. Therefore, a new plasmid was constructed in which the first ATG was missing and in which the second ATG was changed to AAG by means of PCR mutagenesis, as follows. Primers 5LE(E) 5'-CAATGGAATTCAAGGCAAAACAGCTCAAGGTAAATAATACGGG-3' (SEQ ID NO:104); nt 8332-8374) and 3LE (B) 5'-GTGAATCTAGAATGCCGGATCCGTACGAATGC-3' (SEQ ID NO:105); nt 8847-8870) were used in a PCR reaction using plasmid pGEM-L7a (FIG. 3) as a template. The PCR was carried out by using Pwo DNA polymerase and consisted of 30 cycles of 30 seconds at 94°C, 45 seconds at 60°C, and 60 seconds at 72°C. The resulting DNA fragment was digested with EcoRI and XbaI and cloned in pCIneo between the EcoRI and XbaI sites, generating plasmid pCIneoL(N). Subsequently, the BsiWI-SalI fragment from pGEM-L7a, which contains the remaining part of the L gene (nt 8852-15046), was cloned in pCIneoL(N) between the BsiWI and SalI sites, generating plasmid pCIneoL(c). Since antibodies against the L-protein are not available, expression of L could not be checked by immunochemistry.

#### Introduction of a genetic tag in the F gene

**[00149]** To show unambiguously that infectious virus can be generated from cloned full-length cDNA, a genetic tag was introduced in the F gene by means of PCR mutagenesis. To this end, the F gene was cloned by using two overlapping PCR fragments. The first PCR fragment was generated by using primers NDV5F (SEQ ID NO:15) (see above) and primer F5R (5'-AAAGCGCCGCTGTCTCCTCCCTCCAGATGTAGTCAC-3' (SEQ ID NO:106); nt

4859-4894). The residues shown in bold are changes which were introduced in the primer in order to change the amino acid sequence of the proteolytic cleavage site between F1 and F2 from that of the NDV LaSota strain (GGRQGR | L) (SEQ ID NO:135) to that of the consensus cleavage site for virulent NDV strains (GRRQRR | F) (SEQ ID NO:102). The second PCR fragment was generated by using primers F3F (5'-GGAGGAGACAGCGGCGCTTTATAG  
GCGCCATTATTGG-3' (SEQ ID NO:107); nt 4875-4911) and IV09 (5'-CTCTGTCGACACAGACTACCAGAACTTTAC-3' (SEQ ID NO:108); nt 6246-6266).

**[00150]** The PCR was performed with Pwo DNA polymerase and consisted of 25 cycles of 15 seconds at 94°C, 30 seconds at 55°C, and 2 minutes at 72°C. The two overlapping PCR fragments (the overlap is shown in italics in the primer sequences) were joined in a second PCR by using primers NDV5F (SEQ ID NO:15) and IV09 (SEQ ID NO:108) and by using the same PCR conditions. The resulting fragment, which contains the entire ORF of the F gene and which encodes a virulent consensus cleavage site, was digested with NheI and SalI and cloned in pCIneo between the NheI and SalI sites, yielding pCIneoF<sup>wt</sup>. The StuI-NotI fragment (nt 4646-4952) from pCIneoF<sup>wt</sup> was used to replace the corresponding fragment in plasmid p535-S which had been constructed by inserting the ClaI-ScaI (nt 3521-10311) from pGEM-B in p535-DI between the ClaI and ScaI sites (see portion C of FIG. 3). The resulting plasmid was designated p535-S[F<sup>wt</sup>c]. A PCR fragment containing the Cm-resistance gene from pACYC184 (see above) was cloned as an XbaI fragment into the unique XbaI site (position 6172 in the NDV sequence) of plasmid p535-S[F<sup>wt</sup>c], yielding plasmid p535-S[F<sup>wt</sup>c] Cm. Subsequently, the Cm-tagged ApaI-SpeI fragment (nt 2285-8094) of this plasmid was used to replace the corresponding fragment of the full-length cDNA clone pNDFL+. Finally, the Cm gene was removed from this plasmid by digestion with XbaI followed by recircularization using T4 DNA ligase. The resulting plasmid, which contains the genetically tagged full-length NDV cDNA, was designated pNDFL+[F<sup>wt</sup>].

#### Generation of stably transformed cell lines that express individual NDV genes

**[00151]** Plasmids pCIneoNP, pCIneoP, pCIneoM, pCIneoF, pCIneoF<sup>wt</sup>, and pCIneoHN were used for the generation of stably transformed cell lines that express these proteins

individually. The day before transfection, CER cells were seeded in 6 cm culture dishes and incubated overnight to give a confluency of 60-80%. The cells were transfected with 2  $\mu$ g of plasmid DNA by using 12  $\mu$ l of Lipofectamine and OptiMem essentially as described by the supplier (GibcoBRL/Life Technologies). After 48 hours the cells were trypsinized and dilutions were seeded in 10 cm culture dishes in medium containing 500  $\mu$ g/ml of G418 (Boehringer Mannheim). Every 3 days the medium was replaced by fresh medium containing increasing (in steps of 100  $\mu$ g/ml) amounts of G418 until a concentration of 800  $\mu$ g/ml was reached. Cells were kept in medium containing 800  $\mu$ g/ml G418 and three weeks after transfection individual colonies were picked and transferred to 96-well culture dishes. The cloned cell lines were examined for the expression of the respective NDV gene by using an IPMA as described above for transient-expression studies.

**[00152]** Cell lines that constitutively expressed NP, P, M, or F could be identified and isolated. We were unable, however, to generate cell lines that expressed the HN protein. Perhaps constitutive expression of HN is toxic to the cells.

#### Generation of stably transformed cell lines that express T7 RNA polymerase

**[00153]** The gene encoding T7 RNA polymerase was recovered from plasmid pRT7NT (René van Gennip, ID-DLO, Department of Mammalian Virology) by digestion with EcoRI and SalI. The resulting fragment contains the T7 RNA polymerase gene located behind the baculovirus p10 promoter. The DNA fragment was cloned in plasmid pCIneo0 between the EcoRI and SalI sites, generating plasmid pCIneo107. Plasmid pCIneo0 lacks the T7 promoter and was derived from pCIneo by cleavage with NheI followed by partial cleavage with ScaI, filling in the sticky ends with Klenow DNA polymerase and recircularization by using T4 DNA ligase. The baculovirus sequences were removed from pCIneo107 by digestion with EcoRI and PacI, followed by T4 DNA polymerase treatment to generate blunt ends and recircularization. The resulting plasmid was designated pCIneo007. Expression of T7 DNA polymerase was verified by co-transfection of cells with pCIneo007 and pPRh01. The latter plasmid contains the E2 protein of classical swine fever virus cloned behind a T7 promoter and containing an internal ribosome entry site (René van Gennip, personal communication). Expression of E2 was

determined in an IPMA by using monoclonal antibody V4 (Wensvoort et al., 1986). Stably transformed CER cell lines expressing T7 RNA polymerase were generated and isolated as described above except that 10 cm culture dishes were used and the cells were transfected with 5  $\mu$ g of pCIneo007 DNA and 25  $\mu$ l of LipofectAMINE. To examine individual cell lines for the expression T7 RNA polymerase, they were transfected with plasmid pPRh01 and expression of E2 (which is dependent on T7 RNA polymerase) was determined in an IPMA by using monoclonal antibody V4. Several cell lines which expressed T7 RNA polymerase were identified. One cell line, designated CER-C9, was used for subsequent experiments.

#### Cloning and expression of HN genes and hybrid HN genes

**[00154]** Primer 3UIT (SEQ ID NO:1) was used to synthesize single-stranded cDNA of NDV and avian-paramyxovirus serotype-2 and -4 (APMV2 and APMV4) as described above. All subsequent PCR reactions were performed by using 25 cycles of 15 seconds at 94°C, 30 seconds at 55°C and 2 minutes at 72°C.

**[00155]** The entire coding region of the HN gene of APMV2 was recovered by means of PCR by using primers IV03 (5'-GGGGGAATTCCCCATTCAATGAAGGGTCTAC-3' (SEQ ID NO:110)) and IV05 (5'-GATCCCCGGGTCTTAAACCAGGCTTCGCAATG-3' (SEQ ID NO:111)) which were derived from the sequence of the HN gene of APMV2 (GenBank accession number D14030). The entire coding region of the HN gene of APMV4 was recovered by means of PCR by using primers IV06 (5'-GGGGGAATTCTGGTAGGGTGGGGAA GGTAGC-3' (SEQ ID NO:112)) and IV08 (5'-ATTGCCCGGGGGGTAACTAATCAGGATCT CAG-3' (SEQ ID NO:113)) which were derived from the sequence of the HN gene of APMV4 (GenBank accession number D14031). The resulting PCR fragments were digested (either directly or after subcloning in pGEM-T), with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmids were designated pCIneoHN2 and pCIneoHN4, respectively.

**[00156]** Hybrids between the HN gene of NDV strain LaSota and the HN genes of APMV2 and -4 were constructed by means of overlap PCR as follows. The N-terminal part (aa 1-141) of the HN gene of NDV strain LaSota was amplified with Pwo DNA polymerase by using



primers IV01B (5'-GTAGGAATTCAAGAGAGGCCGCCCTCAAT-3' (SEQ ID NO:114); nt 6325-6354) and IV10 (5'-AATGAGTTCTTTGCCTATCCCCC-3' (SEQ ID NO:115); nt 6811-6834). The C-terminal part of the HN gene of APMV2 (aa 142-580) was amplified with Pwo DNA polymerase by using primers IV11B (5'-GGGGGGATAGGCAAAGA ACTCATTCAAGGAGATGCATCTGCAGGC-3' (SEQ ID NO:116) and IV05 (SEQ ID NO:111). The resulting PCR fragments were joined in an overlap PCR (overlap shown in italics) by using primers IV01B (SEQ ID NO:114) and IV05 (SEQ ID NO:111) and by using the Expand High Fidelity enzyme mix. The resulting PCR fragment was digested (either directly or after subcloning in pGEM-T) with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmid which contains a hybrid HN gene consisting of aa 1-141 of NDV and aa 142-580 of APMV2 was designated pCIneoHN1/2<sup>141</sup>.

**[00157]** The C-terminal part of the HN gene of APMV4 (aa 143-569) was amplified by using primers IV14B (5'-GGGGGGATAGGCAAAGAACTCATTGTAGATGATGCATCTGC AGGCCTAAATTTCC-3' (SEQ ID NO:117) and IV08 (SEQ ID NO:113). This fragment was joined with the N-terminal part of the HN gene of NDV (see above) in an overlap PCR by using primers IV01B (SEQ ID NO:114) and IV08 (SEQ ID NO:113). The resulting PCR fragment was digested (either directly or after subcloning in pGEM-T) with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmid which contains a hybrid HN gene consisting of aa 1-141 of NDV and aa 143-569 of APMV4 was designated pCIneoHN1/4<sup>141</sup>.

**[00158]** In analogy to the constructions described above, hybrid HN genes were constructed which consisted of aa 1-143 of NDV and aa 144-580 of APMV2, or aa 1-143 of NDV and aa 145-569 of APMV4. For these constructions PCR fragments were obtained by using the following pairs of primers; NDV aa 1-143, primer IV01B (SEQ ID NO:114) and IV13 (5'-ATCTACAATGAGTTCTTTGCCTATC-3' (SEQ ID NO:118); nt 6816-6840); APMV2 aa 144-580, primer IV14B (5'-GGGGGGATAGGCAAAGAACTCATTGTAGATGATGCATCTGCA GGCCTAAATTTCC-3' (SEQ ID NO:117) and IV05 (SEQ ID NO:111); APMV4 (SEQ ID NO:128) aa 145-569, primer IV15B (5'-GGGGGGATAGGCAAAGAACTCATTGTAGAT CAAACAGCTGACTACACAGCAG-3' (SEQ ID NO:119) and IV08 (SEQ ID NO:113). The

PCR fragments were digested (either directly or after subcloning in pGEM-T) with EcoRI and XmaI and cloned in pCIneo between the EcoRI and XmaI sites. The resulting plasmids were designated pCIneo1/2<sup>143</sup> and pCIneoO1/4<sup>143</sup>, respectively. To examine expression of the HN proteins, CER cells or QM5 cells were infected with FPV-T7 for 1 hour at a m.o.i. of 1, transfected with plasmids pCIneoHN, pCIneoHN2, pCIneoHN4, pCIneoHN1/2<sup>141</sup>, pCIneoHN1/2<sup>143</sup>, pCIneoHN1/4<sup>143</sup> and pCIneoHN1/4<sup>143</sup> and 24 hours after transfection the monolayers were overlaid with a 1% suspension of chicken erythrocytes in PBS for 45 minutes at room temperature. Subsequently, the monolayers were carefully washed three times with PBS and adhesion of erythrocytes to transfected cells was examined microscopically. To examine induction of cell fusion after co-expression of the HN and F protein, CER cells or QM5 cells were co-transfected with pCIneoF<sup>wt</sup> together with either pCIneo-HN1 pCIneoHN2, pCIneoHN4, pCIneoHN1/2<sup>141</sup>, pCIneoHN1/4<sup>141</sup>, pCIneoHN1/2<sup>143</sup> or pCIneoHN1/4<sup>143</sup>. After incubation for 2 to 3 days, the monolayers were washed with PBS, stained for 15 minutes with a Giemsa solution (1:30 dilution in water), and examined microscopically.

#### Cloning of hybrid HN genes in full length genomic NDV cDNA

**[00159]** A synthetic linker, designated HN12, was inserted between the NotI and SpeI sites of pGEM-T (Promega) by using oligonucleotides HN12a (5'-GGCCGCATATTCTAGAGTTAACGACTTA-3' (SEQ ID NO:120) and HN12b (5'-CTAGTAAGTCGTTAAGTCTAGAAATATGC-3' (SEQ ID NO:121)). A synthetic linker, designated HN14, was inserted between the NotI and SpeI sites of pGEM-T by using oligonucleotides HN14a (5'-GGCCGCATATTCTAGAGTTAACGA-3' (SEQ ID NO:122) and HN14b (5'-CTAGTCGTTAAGTCTAGAAATATGC-3' (SEQ ID NO:123). The resulting plasmids were designated pGEM-HN12 and pGEM-HN14, respectively. These plasmids were digested with NotI and XbaI and used to clone the NotI-SpeI fragment (nt 3390-7488) from plasmid p535-S[F<sup>wt</sup>c]Cm. The resulting plasmids were designated pGEM-HN1/2NS and pGEM-HN1/4NS, respectively. The HN genes of these plasmids were replaced by the hybrid HN genes from plasmids pCIneoHN1/2<sup>143</sup> and pCIneoHN1/4<sup>143</sup>, respectively (see section: Cloning and expression of HN genes and hybrid HN genes). To this end, pCIneoHN1/2<sup>143</sup> and

pCIneoHN1/4<sup>143</sup> were digested with NheI and SmaI and the resulting fragments (containing the hybrid HN1/2<sup>143</sup> and hybrid HN1/4<sup>143</sup> genes) were cloned between the NheI and HpaI site of plasmids pGEM-HN1/2NS and pGEM-HN1/4NS, resulting in pGEM+HN12 and pGEM+HN14, respectively. The latter plasmids were used to introduce the hybrid HN genes into the full length genomic cDNA clone of NDV. To this end, plasmids pGEM+HN12 and pGEM+HN14 were digested with NotI and SpeI and the fragment containing either the HN12 or HN14 gene was used to replace the corresponding fragment of pNDFL+, yielding pNDFL+HN1/2<sup>143</sup> Cm and pNDFL+HN1/4<sup>143</sup> Cm, respectively. The Cm gene was removed from these plasmids by digestion with XbaI followed by recircularization using T4 DNA ligase. In order to comply with the “rule-of-six,” a linker was inserted into the unique SpeI site of these plasmids by using self-complementary oligonucleotides. Linker H2 (5'-CTAGCGAGCGCTCG-3' (SEQ ID NO:124) was inserted in plasmid pNDFL+HN1/2<sup>143</sup> and linker H3 (5'-CTAGCGAGCGWGCTCG-3' (SEQ ID NO:125) was inserted in pNDFL+HN1/4<sup>143</sup>, yielding plasmids pNDFL+HN1/2<sup>143</sup> (H2) and pNDFL+HN1/4<sup>143</sup> (H3), respectively.

#### Elimination of a specific epitope in the HN protein of NDV LaSota

[00160] A specific epitope, *i.e.*, amino acids 346 to 354 (PDEQDYQIR) (SEQ ID NO:126), in the HN protein of NDV LaSota that is recognized by MAb 4DE (Long et al., 1986; Meulemans et al., 1986), was eliminated by replacing this sequence by the corresponding sequence of the HN proteins of either APMV-2 (NRTDIQQTI) (SEQ ID NO:127) or APMV-4 (PDPLQDQIL) (SEQ ID NO:128). To this end, plasmid pCIneoHN (see section: Cloning and expression of individual NDV genes) was used as template to create overlapping PCR fragments. For the APMV-2 (SEQ ID NO:127) sequence the first PCR fragment was generated by using primers IV01 (5'-GTAGACGCGTAAGAGAGGCCGCCCTCAAT-3' (SEQ ID NO:129) and primer 3HN2 (5'-GATAGTTTGCTGTATATCAGTCCGATTGCATGTGTCATTGTATCGCT TGTATATCAC-3' (SEQ ID NO:130). The second PCR was generated by using the primers 5HN2 (5'-AATCGGACTGATATACAGCAAACATCATGGCCAAGTCTTCGTATAAGCCT GGAGCC-3' (SEQ ID NO:131) and NDV3-HN (5'-CGAGCCCCGGGCGGCATTCGGT TTGATTCTTG-3' (SEQ ID NO:103)). The resulting fragments were combined and used as

template for a third PCR by using the primers IV01B (5'-GTAGGAATTCAAGAGAGGCCGCCCTCAAT-3' (SEQ ID NO:114)) and primer NDV3-HN (SEQ ID NO:103). For the APMV-4 (SEQ ID NO:128) sequence the first PCR fragment was generated by using primers IV01 (SEQ ID NO:129) and primer 3HN4 (5'-TAAGATCTGATCTTGCAGCGGGTCAGGGCATGTGTCATTGTATCGCTTGTATATC AC-3' (SEQ ID NO:115)). The second PCR was generated by using the primers 5HN4 (5'-CCTGACCGCTGCAAGATCAGATCTTAATGGCCAAGTCTTCGTATAAGCCTGGAGC C-3' (SEQ ID NO:116)) and NDV3-HN (SEQ ID NO:104). The resulting fragments were combined and used as template for a third PCR by using the primers IV01B (SEQ ID NO:114) and NDV3-HN (SEQ ID NO:104). Primers 3HN2/5HN2 and 3HN4/5HN4 are partly complementary and contain the genetic codes for the HN2 sequence (NRTDIQQTI) (SEQ ID NO:127) and HN4 sequence (PDPLQDQIL) (SEQ ID NO:128), respectively. The PCR reactions were performed by using the Expand Long Template PCR kit (Boehringer Mannheim). The PCR consisted of 30 cycles of 10 seconds 94°C, 30 seconds 58°C and 2 minutes at 68°C, followed by 1 cycle of 4 minutes 68°C. The PCR fragments were digested with EcoNI and Bsu36I, and cloned between the EcoNI and Bsu36I sites of pCIneoHN. The resulting plasmids were designated pCIneoHN1(HN2e) and pCIneoHN1(HN4e), respectively. Transient expression studied indicated that the modified HN proteins were correctly expressed and transported to the cell surface as judged from hemadsorption studies using chicken erythrocytes. Furthermore, MAb 6D4 which is directed against a linear epitope of HN of NDV and which consists of (or at least includes) amino acids 346-354, did not react with the modified HN proteins.

**[00161]** Plasmids pCIneoHN1(HN2e) and pCIneoHN1(HN4e) were digested with NarI and SpeI and the fragments containing the modified HN genes were cloned between the NarI and SpeI sites of pGEM-HN1/2NS and pGEM-HN1/4NS, respectively. The resulting plasmids, designated pGEM-HN1(HN2e) and pGEM-HN1(HN4e), were digested with NotI and SpeI, and used to replace the NotI-SpeI fragment in pNDFL+. The resulting plasmids were designated pNDFL-HN(HN2e)Cm and pNDFL-HN(HN4e) Cm, respectively. The Cm gene was removed from these plasmids by digestion with XbaI followed by religation. The resulting plasmids were designated pNDFL-HN(HN2e) and pNDFL-HN(HN4e), respectively.

## RESULTS

Nucleotide sequence of the 3'- and 5'-terminal ends of the genome of NDV strain LaSota

**[00162]** The sequence of a putative 3' end of the NDV genome has been published (Ishida et al., 1986) albeit from another NDV strain (D26) than the one used here (LaSota). Yusoff et al. (1987) have published a sequence of the L gene and a relatively large non-coding region behind the L gene of NDV strain Beaudette C. However, as shown herein, this sequence did not include the full terminal 5' end of the viral genome which makes it impossible to generate infectious copy NDV. The 3'- and 5'-terminal ends of the genome of negative-strand RNA viruses fulfill an essential function in replication and transcription (Lamb and Kolakofsky, 1996). Thus, in order to generate a full-length NDV cDNA which can be used to generate infectious virus by means of reverse genetics (Conzelmann, 1996), it is absolutely essential to include the correct 3' and 5' ends of the viral genome. Therefore, we determined the exact nucleotide sequence of both the 3' and 5' ends of the genomic RNA of NDV strain LaSota by using 3'- and 5'-RACE procedures (rapid amplification of cDNA ends). The 5' end was recovered by means of PCR after ligation of a single-stranded anchor primer (ALG3) (SEQ ID NO:83) to single-stranded cDNA which was generated by reverse transcription of the 5' end of the genomic RNA. By using a primer (ALG4) (SEQ ID NO:85) that is complementary to the anchor primer and an NDV-specific primer, PCR products were generated which contained the 5' end.

**[00163]** To clone the 3' end of NDV, the single-stranded anchor primer ALG3 (SEQ ID NO:83) was ligated to the 3' end of viral RNA by using T4 RNA ligase and amplified by means of PCR by using primer ALG4 (SEQ ID NO:128) and an NDV-specific primer (method I). Alternatively, the 3' and 5' ends of the NDV RNA were ligated to each other by using T4 RNA ligase and the resulting concatenated RNA was used for RT-PCR by using NDV-specific primers that flanked the ligation point (method II). The 3'- and 5'-RACE products were cloned in T-vector pBluescriptII-TSK (Ichihara and Kurosawa, 1993) or in pGEM4Z and several independent clones were isolated and sequenced. The results are compiled in Table 2. To enable the direct comparison of the 3'- and 5'-terminal ends, the sequences are shown in the form of DNA and the 3' end of the genomic strand is represented as the 5' end of the antigenomic strand.

At the genomic RNA level the sequence of the 3' end reads 3'-UGGUUUGUCUCUUAG (SEQ ID NO:132) whereas the sequence of the 5' end reads UUUAGAAACAAACCA-5' (SEQ ID NO:133). The sequence of the 3' end is almost similar to the published 3'-terminal sequence of NDV strain D26 (Ishida et al., 1986). However, the sequence of the 5' end showed that NDV strain LaSota contains 64 additional nucleotides in comparison with the published sequence of the L gene of NDV strain Beaudette C (Yusoff et al., 1987) (FIG. 5.)

#### Replication of NDV minigenomes by helper virus

**[00164]** To determine whether the 3' and 5' ends of NDV are functional in replication and transcription, minigenomes were constructed which consisted of the 3' end of NDV (nt 1-119), a reporter gene encoding secreted alkaline phosphatase (SEAP), and the 5' end of NDV (nt 14973-15186) (FIG. 2). These minigenomes were cloned in both orientations in transcription vector pOLTV5, generating plasmids pOLTV535 and pOLTV553, respectively (for details of the construction see Materials and Methods). Plasmid pOLTV5 (FIG. 1) contains the T7 RNA polymerase promoter followed by unique *Stu*I and *Sma*I restriction sites, the autocatalytic ribozyme from hepatitis delta virus (HDV) and the transcription termination signal from bacteriophage T7 (Pattnaik et al., 1992). *In vivo* or *in vitro* transcription using T7 RNA polymerase of plasmid pOLTV535 gives rise to antigenomic RNA (or [(+)-RNA]), whereas transcription of plasmid pOLTV553 gives rise to genomic RNA (or [(-)-RNA]) (FIG. 4).

**[00165]** To examine whether the minigenome RNA's generated by plasmids pOLTV535 and pOLTV553 could be replicated and expressed by using NDV as helper virus, we used CER cells which expressed T7 RNA polymerase either constitutively (CER-C9 cells, see Materials and Methods), or after infection with fowl pox recombinant fpEFLT7pol (Britton et al., 1995; hereafter called FPV-T7) that expresses T7 RNA polymerase. CER-C9 cells and FPV-T7 infected CER cells were transfected with the minigenome plasmids pOLTV535 or pOLTV553 and after incubation for 3 hours at 37°C the cells were either infected with NDV for 1 hour, or left uninfected. Approximately 24 hours after transfection, a sample was taken from the medium and assayed for SEAP activity. The results showed the SEAP expression as very high in FPV-T7 infected cells which were transfected with pOLTV535. This is not surprising since

transcription by T7 RNA polymerase generates antigenomic [+] RNA which is capped by fowl pox enzymes and which is efficiently translated by the host cell. In cells transfected with pOLTV553, transcription by T7 RNA polymerase generates genomic [-] RNA which must be converted, into [+] RNA by helper virus in order to be translated into SEAP protein (cf. FIG. 4). In both cases, no increase in SEAP expression could be observed in NDV infected cells in comparison to non-infected cells. On the contrary, SEAP-expression in NDV infected cells was consistently approximately two times lower than in uninfected cells (results not shown). For pOLTV535-transfected cells this may be explained by the already very high level of SEAP expression by transcripts generated by T7 RNA polymerase. However, in pOLTV553-transfected cells, where efficient expression of SEAP is dependent on the conversion of genomic [-] RNA into antigenomic [+] RNA or mRNA by the viral polymerase complex, we would have expected an increase in SEAP expression after NDV infection.

**[00166]** We could think of two reasons why the minigenomes could not be expressed and replicated by NDV. First, the size of the minigenome RNA's does not conform to the so-called "rule-of-six" (Calain and Roux, 1993; Kolakofsky et al., 1998). According to this rule, paramyxovirus genomes are only replicated efficiently when they are a multiple 6 nt in length. Second, the two extra G residues which are present at the 5' end of the minigenome RNA's might interfere with correct replication and/or transcription by the viral polymerase complex. To find out whether replication of the genomes was dependent on the rule-of-six, we inserted a series of short self-complementary oligonucleotides which increased 1 nt in size in the unique ClaI site in plasmids pOLTV535 and pOLTV553 (FIG. 2). The resulting plasmids (pOLTV535N0 to -N5 and pOLTV553N0 to -N5) differ in size from 1 to 6 nt and therefore one of them should generate a minigenome RNA which conforms to the rule-of-six. The plasmids were used to transfect CER cells or FPV-T7 infected CER-C9 cells as described above. The results showed that only plasmids pOLTV535N3 and pOLTV553N3 gave rise to an enhanced SEAP activity after NDV infection. The length of the minigenome RNA's generated from these plasmids by T7 RNA polymerase were calculated to be  $6n+2$ . Since two extra G residues are present at the 5' end of the minigenome RNA's, these results suggest that only the size of the RNA sequence which is located between the authentic 3' and 5' ends of the minigenome RNA's

is relevant for the rule-of-six. This was verified by constructing minigenome plasmids in which the transcription start of T7 RNA polymerase was changed so that the first nucleotide which was incorporated into RNA was the first nucleotide of the 3' or 5' end of NDV (see Materials and Methods). Transfection of these plasmids indicated that only minigenome RNA's generated by plasmids pOLTV735N3 and pOLTV753N3 are replicated by helper virus (results not shown). These findings again indicate that replication of NDV is strictly dependent on the rule-of-six. Furthermore, these findings indicate that the presence of two extra G residues at the 5' end of the minigenome RNA's does not interfere with correct replication. Similar results have been obtained with minigenome plasmids (or DI plasmids) from other paramyxoviridae (Pattnaik et al., 1992; Harty and Palese, 1995).

#### Packaging of NDV minigenomes by helper virus

**[00167]** To determine whether minigenome RNA's could be packaged by NDV helper virus, the medium of the transfected cells was transferred to fresh monolayers and after 1 hour of adsorption, the monolayers were washed three times with PBS and further incubated in complete medium. After 24 hours of incubation, the SEAP activity in the medium was measured. The results showed that SEAP activity was present only in cells which had been treated with the medium from cells transfected with minigenome plasmid pOLTV553N3 (Table 4). This finding indicates that minigenome RNA's can be packaged into NDV envelopes and that these particles are able to infect cells. Furthermore, these results show that packaging is dependent on replication which indicates that only RNA molecules which are complexed with the viral NP, P and L proteins are packaged into virus-like particles.

#### Replication of NDV minigenomes by plasmids expressing the NP, P, and L proteins

**[00168]** To determine whether the minigenome RNA's could also be replicated by plasmids encoding the essential NP, P, and L proteins, we performed co-transfection experiments in cells infected with FPV-T7. Cells were transfected with a combination of plasmids consisting of the minigenome plasmid and plasmids pCIneoNP, -P, and -L(c), respectively. As a negative control, pCIneoL(c), which encodes the essential L protein, was replaced by the vector plasmid



pCIneo. The results (Table 5) indicated that indeed plasmids encoding NP, P, and L are able to replicate minigenome RNA's. The results furthermore show that, similar to minigenome replication by helper virus, also replication by the NP, P, and L proteins is dependent on the rule-of-six.

#### Nucleotide sequence of the complete genome of NDV strain LaSota

**[00169]** Sub-genomic cDNA fragments spanning the entire NDV genome were constructed by means of RT-PCR (FIG. 3). To keep the number of PCR errors to a minimum, a proofreading enzyme-mix (Expand Long Template; Boehringer Mannheim) was used in combination with a limited number of PCR cycles (15 cycles). Primer 3UIT (SEQ ID NO:1) which is complementary to the 3' end of NDV RNA was used for reverse transcription, and gene-specific primers were used for PCR. To identify possible PCR errors, three independent RT reactions were performed and used to generate three independent sets of subgenomic cDNA's. The cDNA's, which varied in size from approximately 4 to 7 kb, were cloned in pGEM-T. The nucleotide sequence of two sets of cDNA's was determined by using primers which were either deduced from published NDV sequences, or by primers derived from the NDV sequence that was deduced during this sequencing project (Table 1). Remaining ambiguities were resolved by sequencing the relevant regions of the third set of cDNA clones. The genome of NDV strain LaSota consists of 15186 nt (SEQ ID NO:134), which makes it the smallest of all paramyxovirus genomes from which the entire sequence has been established to date (Kolakofsky et al., 1998).

#### Construction of a full-length NDV cDNA clone in transcription plasmid pOLTV5

**[00170]** To construct a full-length cDNA clone of NDV strain LaSota, overlapping cDNA clones spanning the entire NDV genome were joined at shared restriction sites according to the strategy shown in FIG. 3. The entire NDV cDNA was assembled in the minigenome plasmid pOLTV535 (see above) which is derived from transcription plasmid pOLTV5.

**[00171]** As can be seen in portion B of FIG. 3, the last step in the assembly of the complete NDV cDNA was the cloning of an approximately 8.8 kb *Cla*I (nt 3521-12355)

fragment from pGEM-B into p535-DI which contained the NDV sequences flanking the ClaI site at either side (i.e., nt 1-3521 and 12355-15186, respectively). This step proved to be quite difficult since we repeatedly failed in generating the correct clones. Therefore, the ClaI fragment of pGEM-B was tagged with the chloramphenicol-resistance (Cm) gene from plasmid pACYC184. The ClaI fragment harboring the Cm gene was isolated and cloned in the ClaI site of p535-DI and transformants were selected for resistance against both Cm. Since transformants grew poorly, the antibiotic selection was reduced to 15 µg/ml Cm and 10 µg/ml Km and the incubation temperature was reduced from 37°C to 32°C. Finally, the Cm gene was removed from this plasmid by digestion with BsiWI followed by recircularization by using T4 DNA ligase. After transformation of *E. coli*, cells harboring the desired plasmid were identified phenotypically by screening for Km-resistance and Cm-sensitivity. The resulting plasmid which consisted of the full-length NDV cDNA cloned between the SmaI and StuI sites of transcription plasmid pOLTV5 was designated pNDFL+.

#### Generation of infectious NDV from full-length cDNA

**[00172]** To generate infectious NDV entirely from cloned cDNA, plasmid pNDFL+ was used in co-transfection experiments with pCIneoNP, -P, and -L(c), as described above for the minigenome plasmids. Transfection of CER and CEF cells was monitored by using minigenome plasmid pOLTV553N3 and by measuring SEAP expression. As a negative control, pCIneoL(c) was replaced by pCIneo. After co-transfection, the cells were incubated for 3 to 6 days in medium containing 5% allantoic fluid. The addition of allantoic fluid is necessary because CER or CEF cells lack the appropriate proteases which are required to cleave the F protein of NDV strain LaSota. Cleavage of the F protein is absolutely required for cell-to-cell spread and for the generation of infectious virus. After 3 days of incubation, we performed an immunological staining of the fixed monolayers by using a monoclonal antibody against the F protein. The results showed that cells that were stained with the antibody were only present in monolayers which had been co-transfected with pNDFL(+), pCIneoNP, -P, and -L(c). These results indicated that genome replication and expression was occurring in these cells. No staining cells were observed when pCIneoL(c) was replaced by pCIneo in the co-transfection experiments.

**[00173]** To recover infectious virus, the supernatant of transfected CEF monolayers was injected into the allantoic cavity of embryonated eggs. Four days later the allantoic fluid was harvested, analyzed in a hemagglutination assay, and passaged further in eggs. The results showed that only the supernatant of cells transfected with a combination of pNDFL+ and pCIneoNP, -P, and -L(c) yielded a positive reaction in the hemagglutination assay. Allantoic fluid which showed a positive hemagglutination reaction was subsequently analyzed in a hemagglutination-inhibition assay by using monoclonal antibodies 7B7, 8C11, 5A1, 7D4, and 4D6 (Long et al., 1986) which can be used to differentiate between different NDV strains. The results of this assay indicated that the NDV strain which was recovered from the inoculated eggs showed the same reactivity as the original LaSota strain. The virus which was recovered from the inoculated eggs was designated NDFL to distinguish it from the original LaSota strain.

#### Generation of genetically modified NDV from full-length cDNA

**[00174]** To show unambiguously that the co-transfection system could be used to recover infectious virus from cloned full-length NDV cDNA, a genetic tag was introduced in plasmid pNDFL(+). To this end, the amino acid sequence of the protease cleavage site in the F<sub>0</sub> protein was changed from that of the LaSota strain (GGRQGR | L) (SEQ ID NO:134) to the consensus sequence of virulent NDV strains (GRRQRR | F) (SEQ ID NO:135) by means of PCR mutagenesis (for details see Materials and Methods). The resulting plasmid, pNDFL+[F<sup>wt</sup>], was used to generate virus by using the co-transfection system described above. Infectious virus, designated NDFL[F<sup>wt</sup>], was recovered from the allantoic fluid of embryonated eggs which had been inoculated with the medium of co-transfected CEF cells. In an HI test, all Mabs including 7D4, which is specific for the LaSota strain, showed the same reactivity with the newly generated virus as with the original LaSota strain. The nucleotide sequence of the region encoding the protease cleavage site of the F protein was determined by means of RT-PCR. The results showed that the nucleotide sequence contained the exact nucleotide changes which were introduced in the mutagenic primer which was used to modify the original LaSota sequence. This finding shows that the virus was derived from plasmid pNDFL+[F<sup>wt</sup>] and demonstrates that (genetically modified) NDV can be generated entirely from cloned full-length NDV cDNA.

The protease cleavage site of the Fo protein of NDV is a key determinant for virulence

[00175] It is generally assumed that the amino acid sequence of the protease cleavage site of the Fo protein is a key determinant for virulence of different NDV strains. The generation of a genetically modified LaSota strain in which the amino acid sequence of the protease cleavage site was changed from a lentogenic (non-virulent) to that of a velogenic (virulent) NDV strain offered the unique opportunity to test this assumption. Therefore, we determined the intracerebral pathogenicity index (ICPI) of the newly generated virus NDFL[F<sup>wt</sup>] and compared it with that of strain NDFL and of the original LaSota strain (clone E13-1). The results showed that the ICPI of strain NDFL[F<sup>wt</sup>] was 1.3 which is far above the value for strains NDFL (ICPI=0.0) and clone E13-1 (ICPI=0.3). These results show that, as expected, the virulence of NDV is largely determined by the amino acid sequence of the protease cleavage site of the Fo protein.

#### Introduction of serological marker

[00176] The envelope glycoproteins F and HN of NDV are the most immunogenic proteins of the virus. After infection, both the F and HN protein elicit a strong neutralizing antibody response. The induction of such a neutralizing antibody response is the basis of successful vaccination by non-virulent NDV strains (such as the widely used LaSota strain). However, the antibody response against NDV vaccine strains cannot be distinguished from the antibody response against virulent NDV field strains. Thus, infections with virulent field virus cannot be traced by serological methods. This situation is undesirable since field virus infections are masked by vaccination and clinical signs which are caused by field strains may be overlooked or are even attributed to the vaccine. Since successful differentiation between vaccination and infection is essential for eradication of NDV, we set out to develop genetically modified NDV strains which can be used for vaccination and which can be serologically distinguished from NDV field strains (so called marker vaccines).

[00177] In order to develop an NDV marker vaccine, the virus has to be genetically modified such that one or several immunodominant epitopes of one of the (major) antigens are

either deleted or modified. Deletion of part(s) of an essential protein may lead to the loss of the biological function of that protein. Therefore, we chose to modify one of the immunodominant envelope proteins of NDV in such a way that the biological function of the protein was retained whereas the antibody repertoire against the modified protein differed from that against the original protein. For reasons specified below, we chose for one embodiment of the invention to modify the HN protein of NDV. Infection of NDV is initiated by fusion of the virion envelope with the plasma membrane of the host cell. For this process, both the F protein and the HN protein are required. It has been shown that the F and HN proteins physically interact and that this interaction is required for membrane fusion (Deng et al., 1995). Furthermore, it has been shown that the interaction is type specific, i.e., the F and HN proteins must be derived from the same virus in order to show fusion activity. The interacting domain of the HN protein of NDV has been localized to the so-called stalk- or stem-region of the protein, comprising the first 92 amino acid residues of the ectodomain of the HN protein (Deng et al., 1995). Hybrid HN proteins consisting of aa 1-141 of NDV and aa 141-572 of human parainfluenza virus type-3 (hPIV3) were shown to retain fusion activity when co-expressed with the NDV F protein. These findings suggest that genetically modified NDV strains which harbor a hybrid HN protein which consists of the stem region of NDV followed by the globular head of the HN protein of a different avian-paramyxovirus serotype may be viable. Furthermore, such strains would elicit an anti-HN antibody response which is different from that of NDV. Since the neutralizing antibody response against the F protein is sufficient to allow efficient protection against challenge virus infection, such genetically modified NDV strains meet the two essential requirements of a marker vaccine, i.e., protection against disease and serological differentiation.

**[00178]** Hybrid HN genes were constructed which consisted of a fusion of either aa 1-141 of NDV and aa 142-580 of avian-paramyxovirus type-2 (APMV2) (designated HN1/2<sup>141</sup>) or aa 1-143 of NDV and aa 144-580 of APMV2 (designated HN1/2<sup>143</sup>). Similarly, hybrid HN genes were constructed which consisted either of aa 1-141 of NDV and aa 143-569 of APMV4 (designated HN1/4<sup>141</sup>) or aa 1-143 of NDV and aa 145-569 of APMV4 (designated HN1/4<sup>143</sup>). The hybrid genes were cloned in the eukaryotic-expression vector pCIneo and used in co-transfection experiments with a plasmid harboring the NDV F protein. To this end, the F

protein was modified such that the amino acid sequence of the proteolytic cleavage site between F2 and F1 was changed from the LaSota sequence to that of the consensus sequence of virulent NDV strains ( $F^{wt}$ , see Materials and Methods section). Co-transfection experiments in CER cells and QM5 cells indicated that both HN1/2<sup>141</sup> and HN1/2<sup>143</sup> as well as HN1/4<sup>141</sup> and HN1/4<sup>143</sup> induced cell fusion when co-expressed with the  $F^{wt}$  protein. These results indicated that the complexes between the hybrid HN proteins and the F protein were biologically active. The hybrid HN proteins HN1/2<sup>143</sup> and HN1/4<sup>143</sup> were used to replace the original HN gene in the full-length cDNA clone pNDFL+, yielding pNDFL-HN1/2<sup>143</sup> and pNDFL-HN1/4<sup>143</sup>. The latter two plasmids were subsequently used for the generation of infectious virus by using the co-transfection system described above. Viable recombinant viruses (designated NDFL-HN1/2<sup>143</sup> and NDFL-HN1/4<sup>143</sup>) could be isolated from the allantoic fluid of embryonated eggs which had been inoculated with the supernatant of transfected monolayers.

**[00179]** The presence of the hybrid HN gene in each of two recombinants was verified by means of RT-PCR. Hemagglutination-inhibition tests showed that monoclonal antibodies and polyvalent antisera against NDV were unable to inhibit agglutination of chicken erythrocytes by the recombinant viruses NDFL-HN1/2<sup>143</sup> and NDFL-HN1/4<sup>143</sup>. These results indicate that strains NDFL-HN1/2<sup>143</sup> and NDFL-HN1/4<sup>143</sup> may be used as vaccines that can be serologically distinguished from classical NDV vaccines.

#### Expression of a heterologous protein from recombinant NDV

**[00180]** To examine whether foreign genes can be inserted into the NDV genome, we constructed a recombinant virus that carried the SEAP reporter gene. The SEAP gene was derived from plasmid pOLTV535 and was modified to include the typical transcriptional stop and start boxes of NDV. A DNA fragment containing the SEAP gene followed by the transcriptional stop and start boxes was inserted into the XmnI site (nt 109) in plasmid pNDFL+[( $F^{wt}$ )]. Infectious virus, designated NDFL-AP, was generated by means of the co-transfection system, and the presence of the SEAP gene was verified by means of RT-PCR. Cells infected with strain NDFL-AP expressed very high levels of the SEAP protein. By using the specific activity of the SEAP protein, we calculated that x% of the proteins expressed in cells

infected with NDFL-AP consisted of SEAP protein. These results show that heterologous genes can be expressed to very high levels from recombinant NDV.

#### Generation of an NDV deletion mutant on a trans-complementing cell line

**[00181]** In order to abrogate expression of the M protein of NDV, a large part of the M gene was deleted by digestion of pNDFL+[F<sup>wt</sup>] with BsaAI (nt 3087) followed by partial digestion with HindIII (nt 4252). After filling in the HindIII end with Klenow DNA polymerase, the fragment was recircularized by using T4 DNA ligase and used to transform *E. coli*. The resulting plasmid, designated pNDFL+[F<sup>wt</sup>]dM, was used to generate virus by means of the co-transfection system in trans-complementing CER-M cells that expressed the NDV M protein. The supernatant of transfected monolayers was passaged three times on CER-M cells and analyzed for the presence of virus. Virus was obtained as evidenced by the fact that the culture supernatant of the third passage yielded positive results in hemagglutination (HA) and hemagglutination-inhibition (HI) tests. The virus was designated NDFL-dM. When NDFL-dM was used to infect monolayers of CEF cells, the virus was still able to spread by cell-to-cell transmission as seen in an IPMA by using a monoclonal antibody against the F protein. As expected, expression of the M protein could not be demonstrated in an IPMA by using monoclonal antibodies against the M protein. When the supernatant was used to infect either CEF cells or CER-M cells, we were unable to show the presence of replicating virus in these monolayers by means of IPMA. This finding indicates that infectious virus could not be generated in non-complementing CEF cells. This finding was confirmed by the observation that inoculation of embryonated eggs with supernatant from infected CEF cells did not result in the generation of progeny virus when tested in HA or HI tests.

**[00182]** The need for better NDV vaccines, and especially the need for NDV marker vaccines, prompted us to develop a reverse genetics system which would allow the genetic modification of NDV. In this document we describe the generation of infectious NDV entirely from cloned full-length cDNA. We show that the virulence of NDV can be dramatically changed by modifying only 3 nucleotides which determine the specificity of the protease cleavage site of the F protein. In this case the protease cleavage site was changed from that of

the LaSota strain to that of the consensus cleavage site of virulent NDV strains. By generating this genetically modified NDV strain we deliver the formal proof that the cleavability of the F protein is the key determinant (but not the only determinant) for virulence of NDV. By using the same reverse genetics approach, the cleavage site can be modified, at will, to any other amino acid sequence. This may lead to the generation of a series of NDV strains which display a spectrum of virulence levels.

#### *In vivo*

**[00183]** As previously mentioned, it has been shown that, besides the cleavability of the F and HN proteins, other viral factors may contribute to pathogenicity. Alterations in transcription and translation can modulate growth and cell-to-cell spread of the virus and/or cytopathogenicity. The availability of an infectious cDNA of NDV allows for the systematic modification of sequences which are involved in transcription and replication. This may lead to the design of new NDV vaccines which sport optimal immunogenicity to virtually non-existing virulence.

**[00184]** Safety is one of the most important properties of live vaccines. However, for many live vaccines, including NDV, immunogenicity is often inversely related to virulence. Therefore, further attenuation of live vaccines without losing immunogenicity is one of the most desired alterations for which genetic modification could be used. In this respect it is worthwhile mentioning that it has been shown that elimination of expression of the V protein of Sendai virus resulted in a markedly reduced *in vivo* pathogenicity for mice (Kato et al., 1997). Similar to Sendai virus, NDV also generates a V protein by a mechanism known as RNA editing (Steward et al., 1993). It is predictable that elimination of expression of the V protein of NDV may also result in an attenuated phenotype *in vivo*.

**[00185]** Apart from changing the virulence of NDV, we show that it is possible to modify the antigenic make-up of NDV in such a way that strains can be generated which can be serologically discriminated from NDV field strains. These, so called, marker vaccines are an invaluable tool to assess the prevalence of NDV in commercial flocks around the world. Furthermore, the large-scale application of such marker vaccines may ultimately lead to the



complete eradication of NDV by a process of intensive screening and stamping out of infected flocks. In this document we show that foreign genes can be inserted into the genome of NDV. These foreign genes can be expressed to very high levels in infected cells. This shows that NDV can be used as a vaccine vector for the expression of antigens from other (poultry) pathogens. Several properties make NDV an ideal vaccine vector for vaccination against respiratory or intestinal diseases. 1) NDV can be easily cultured to very high titers in embryonated eggs. 2) Mass culture of NDV in embryonated eggs is relatively cheap. 3) NDV vaccines are relatively stable and can be simply administered by mass application methods such as addition to drinking water or by spraying or aerosol formation. 4) The natural route of infection of NDV is by the respiratory and/or intestinal tracts which are also the major natural routes of infection of many other poultry pathogens. 5) NDV can induce local immunity despite the presence of circulating maternal antibody.

**[00186]** Finally, we show that viable NDV deletion mutants can be generated by using trans-complementing cell lines. An NDV deletion mutant was generated which is unable to express the matrix (M) protein which is involved in budding of NDV at the inner cell membrane. We show that a phenotypically complemented NDV strain that is unable to express the M protein is still able to infect cells and spread by means of cell-to-cell transmission. However, the mutant virus is unable to generate infectious progeny on non-complementing cells. This finding shows that phenotypically complemented NDV deletion mutants may be used as safe self-restricted vaccines which are unable to spread into the environment. Such a non-transmissible vaccine combines the most important advantage of live vaccines, *i.e.*, efficacy, with the most important advantage of killed vaccines, *i.e.*, safety.

**[00187]** Table 1. Nucleotide sequence of primers.

3' UIT	ACCAAACAGAGAATCCGTGAGTTA (SEQ ID NO:1)	1-24
P368+	GTGATGAGGAACCATGTTGC (SEQ ID NO:2)	368-387
P800+	GTCCGCATCTTCTTGGTTAG (SEQ ID NO:3)	800-819
P1201+	GAGACTTGGAGTAGAGTACG (SEQ ID NO:4)	1201-1220
P1279+	AGCAGCAATGAAGGGCCTGG (SEQ ID NO:5)	1279-1298
P1356+	AAATCGGAGTCCTCACTGGG (SEQ ID NO:6)	1356-1375
P1683+	CTCTATATGACCACACCCTC (SEQ ID NO:7)	1664-1683
PRT1	CAAAGAATTCAGAAAAAAGTACGGGTAGAAG (SEQ ID NO:8)	1785-1814
P2357+	GGAAACAGTCAGGAAAGACC (SEQ ID NO:9)	2358-2377
P2599+	TAAGTAAAGTTGACTATCAG (SEQ ID NO:10)	2599-2618
P2852+	GGCACTTAATAAACTTTCGC (SEQ ID NO:11)	2852-2871
P3496+	GAATGAAGAAGCCACTGTCTG (SEQ ID NO:12)	3496-3515
P3587+	CGGAGATCTTGTTGAGTTGG (SEQ ID NO:13)	3589-3608
P4267+	CATTATCCAAGCAGGTACCC (SEQ ID NO:14)	4270-4299
NDV5-F	ACGGGCTAGCGATTCTGGATCCCGGTTGG (SEQ ID NO:15)	4498-4526
P4731+ (LS)	AAGCTCCTCCCGAATCTGCC (SEQ ID NO:16)	4733-4752
P4958+	AGCTCTGATACAAGCCAAAC (SEQ ID NO:17)	4960-4979
P5266+ (LS)	CTGGTGGGAATATGGATTAC (SEQ ID NO:18)	5267-5286
P5591+ (LS)	AGTAACGTTCCCTATGTCCC (SEQ ID NO:19)	5593-5612
P5616+	GTATTTATTCCTGCTTGAGC (SEQ ID NO:20)	5616-5635
P6000	AATACCCTTGATCAGATGAGAGCC (SEQ ID NO:21)	6166-6190
NDV5-HN	GTAGGCTAGCAAGAGAGGCCGCCCTCAAT (SEQ ID NO:22)	6325-6354
P6693+ (L)	CATTGTAAAAACTGAGACC (SEQ ID NO:23)	6695-6714
P7110+ (L)	ATCGGAAGTCTTGCAGTGTG (SEQ ID NO:24)	7112-7131
P7501+ (L)	TGGTGGGAAACGCATCCAGC (SEQ ID NO:25)	7503-7522

Table 1 (continued)		
P7900+ (LS)	AAGACTTAATCCTACGTCTG (SEQ ID NO:26)	7902-7921
P8590+	AACTCGGAAGGGCAGTACAC (SEQ ID NO:27)	8592-8611
L9000	TTTGTCACCTCTGAACTTGTCATT (SEQ ID NO:28)	9008-9031
P9359+	CAATGATATAGCAGAATCCG (SEQ ID NO:29)	9361-9380
P9371+	GCAGAATCCGTGACTCATGC (SEQ ID NO:30)	9371-9411
P9390+	ATAGCTACTGTATTCTCTGG (SEQ ID NO:31)	9392-9411
P9686+	TCACACGATATCATGTTGAG (SEQ ID NO:32)	9686-9705
P9799+	CACACCCTAACGATAATTGG (SEQ ID NO:33)	9801-9820
P10198+	ATAAGAAACGTATCACTGAC (SEQ ID NO:34)	10200-10219
P10601+	TTGTCGCGTTGCCTGTATGG (SEQ ID NO:35)	10603-10622
P11006+	GCAGACATACTTTGACTCTG (SEQ ID NO:36)	11008-11027
P11393+	TCCCTTATTGTCTGGAGTGC (SEQ ID NO:37)	11395-11414
P11798+	TGATACGATAGAACTCGTAG (SEQ ID NO:38)	11800-11819
L12000	CATATGTCGCCACATGTGAAGGCT (SEQ ID NO:39)	12008-12031
P12373+	CAACCAGGACATATGATGAG (SEQ ID NO:40)	12375-12394
P12796+	TCGACTGTTCTTACCAACTC (SEQ ID NO:41)	12798-12817
P12978+	CACACCAACTTGCAGATACG (SEQ ID NO:42)	12978-12997
P13236+	GAGTATCTACTGTCGGATGC (SEQ ID NO:43)	13238-13257
P13601+	ATACTTGTTTCAGAGGAATAG (SEQ ID NO:44)	13603-13622
P13943+	GACCTGACCTCAGATAAAGC (SEQ ID NO:45)	13946-13965
P14002+	TATCATTGCTGCATTGTGAC (SEQ ID NO:46)	14004-14023
P360	GGCGATGTAATCAGCCTAGTGCTT (SEQ ID NO:47)	14756-14779
P14812+	ACTAAGGACATACTTGAAGC (SEQ ID NO:48)	14812-14831
P230-	CCGGGACTTCTACTTTTAAG (SEQ ID NO:49)	230-211
P998-	TTTGGATATCGCCTGAGAGG (SEQ ID NO:50)	998-979
P1898-	AAAGGTGGCCATGTTTGTCC (SEQ ID NO:51)	1898-1879

Table 1 (continued)		
P2617-	TGATAGTCAACTTTACTTAC (SEQ ID NO:52)	2617-2598
P3328-	GCAGAATCAAAGTACAGCCC (SEQ ID NO:53)	3330-3311
P3610-	CTTGCCAACTCAACAAGATC (SEQ ID NO:54)	3612-3593
P3990-	GATTAGCATAGTATCCACTG (SEQ ID NO:55)	3992-3973
NDV3-M	TCTCCCCGGGGCAGCTTATTTCTTAAAAGGAT (SEQ ID NO:56)	4400-4368
P4593-	GACAGATGCAACTCAGTACC (SEQ ID NO:57)	4625-4606
P4618- (LS)	ATGCAACTCAGTACCAGCGC (SEQ ID NO:58)	4620-4601
P5390-	GTAGAGTTACCTGTATACCC (SEQ ID NO:59)	5411-5392
NDV3-F	ACTACCCGGGAAACCTTCGTTCTCAT (SEQ ID NO:60)	6238-6212
P6710- (LS)	TCTCAGTTTTTAACAATGCC (SEQ ID NO:61)	6712-6693
P7093- (LS)	GTTGATGGAACGCAGAGTAG (SEQ ID NO:62)	7095-7076
P7522- (LS)	CTGCTGGATGCGTTTCCCAC (SEQ ID NO:63)	7524-7505
P367	AGGGACCTCAATACTAGCCAGTTC (SEQ ID NO:64)	8692-8666
P9905-	CTCTATCAAGAGGCGATTAG (SEQ ID NO:65)	9907-9888
P10320-	TAAGACAGTACTTTTGCAGG (SEQ ID NO:66)	10322-10303
P10684-	GATGCAACTGTGTCAACACC (SEQ ID NO:67)	10687-10706
P11122-	AATTGGGCAGGAGTCAGAAC (SEQ ID NO:68)	11124-11105
P11510-	TGCCTCCATGATAGCATGCG (SEQ ID NO:69)	11512-11493
P11903-	ATTGCTTGGAAGATGGAACC (SEQ ID NO:70)	11905-11886
P12717-	TGTCATACACTATTATGGCG (SEQ ID NO:71)	12719-12700
P13141	CAAAGAGTACCGTGTACAGACAGCATAACC (SEQ ID NO:72)	13172-13143
P13281-	GACATGATAGAGCTCACCTG (SEQ ID NO:73)	13302-13283
P14101-	ACGGAATGCATGGCAATCAG (SEQ ID NO:74)	14163-14144
P14522-	GCTCACCAAACCTCTCTGCAC (SEQ ID NO:75)	14524-14505
P14687-	AGGATCTGTCTCGTGCACTG (SEQ ID NO:76)	14709-14690

Table 1 (continued)		
P377	TTTCCTTAAGTTTGGTAATACCTAGGAC (SEQ ID NO:77)	14888-14861
P359	CACCAAGTCGACAATTGGCCAGAAAAGGAG (SEQ ID NO:78)	15046-15017
5NDV	ACCAAACAAAGATTGGTGAATGACGA (SEQ ID NO:79)	15186-15159

**[00188]** Table 2. Sequence of 3'- and 5'-terminal ends of the genome of NDV strain La Sota

A. Sequence of 3'-terminal end (shown as 5' end of antigenomic DNA strand)

method I.	clone	sequence
	04	ACCAAACAGAGAATC (SEQ ID NO:80)
	05	ACCAAACAGAGAATC (SEQ ID NO:80)
	13	ACCAAACAGAGAATC (SEQ ID NO:80)
	21	ACCAAACAGAGAATC (SEQ ID NO:80)
method II.	clone	sequence
	26	ACCAAACAGAGAATC (SEQ ID NO:80)
	28	ACCAAACAGAGAATC (SEQ ID NO:80)
	30	ACCAAACAGAGAATC (SEQ ID NO:80)
	31	GCCAAACAGAGAATC (SEQ ID NO:81)
	32	ACCAAACAGAGAATC (SEQ ID NO:80)
	33	ACCAAACAGAGAATC (SEQ ID NO:80)
	Consensus	<u>ACCAAACAGAGAATC</u> (SEQ ID NO:80)

B. Sequence of 5'-terminal end (shown as DNA)

pBluescriptII-TSK clones	clone	sequence
	r3101-13	ACCAAACAAAGATTT (SEQ ID NO:82)
	r3101-14	ACCAAACAAAGATTT (SEQ ID NO:82)
	r3101-15	ACCAAACAAAGATTT (SEQ ID NO:82)
	r2601-17	ACCAAACAAAGATTT (SEQ ID NO:82)
	r2601-18	ACCAAACAAAGATTT (SEQ ID NO:82)
	r2601-19	ACCAAACAAAGATTT (SEQ ID NO:82)
	r2601-20	AACAAGGTGAAGATA (SEQ ID NO:149)

Table 2 (continued)

	r2601-21	ACCAAACAAAGATTT (SEQ ID NO:82)
pGEM4Z clones	clone	sequence
	r3101-16	ACCAAACAAAGATTT (SEQ ID NO:82)
	r3101-17	ACCAAACAAAGATTT (SEQ ID NO:82)
	r3101-18	ACCAAACAAAGATTT (SEQ ID NO:82)
	r3101-19	ACCAAACAAAGATTT (SEQ ID NO:82)
	r3101-22	ACCAAACAAAGATTT (SEQ ID NO:82)
	Consensus	<u>ACCAAACAAAGATTT</u> (SEQ ID NO:82)

[00189] Table 3. Minigenome replication by NDV helper virus

A. SEAP activity (cps) after transfection of CER-C9 cells with the pOLTV535 and pOLTV553-series of plasmids.

Plasmid	+NDV	-NDV	ratio
pOLTV535N0	$3.5 \times 10^4$	$7.1 \times 10^4$	0.49
pOLTV535N1	5.9	12.1	0.49
pOLTV535N2	2.4	6.2	0.39
pOLTV535N3	7.6	5.2	1.46
pOLTV535N4	1.8	4.1	0.44
pOLTV535N5	1.5	3.0	0.50
pOLTV553N0	$5.5 \times 10^3$	$9.6 \times 10^3$	0.57
pOLTV553N1	9.6	27.6	0.35
pOLTV553N2	2.4	3.5	0.68
pOLTV553N3	15.1	9.5	1.59
pOLTV553N4	3.4	7.9	0.43
pOLTV553N5	2.9	4.8	0.60

B. SEAP activity (cps) after transfection of FPV-T7 infected CER cells with the pOLTV553-series of plasmids.

Plasmid	+NDV	-NDV	ratio
pOLTV553N0	$7.2 \times 10^4$	$8.3 \times 10^4$	0.86
pOLTV553N1	8.4	12.0	0.70
pOLTV553N2	8.9	12.6	0.71
pOLTV553N3	27.4	8.6	3.19
pOLTV553N4	9.7	10.4	0.93
pOLTV553N5	8.5	8.1	1.05



**[00190]** Table 4.

Transfer of SEAP activity (cps) after treatment of CER cells with the supernatant of FPV-T7 infected CER cells which had been transfected with the pOLTV553-series of plasmids and which had been superinfected with NDV (see Table 3).

Plasmid

pOLTV553N0	$2.4 \times 10^3$
pOLTV553N1	6.2
pOLTV553N2	2.0
pOLTV553N3	20.6
pOLTV553N4	2.0
pOLTV553N5	2.1

**[00191]** Table 5.

SEAP activity (cps) after co-transfection of CER cells with the pOLTV553-series of plasmids and plasmids pCIneoNP, pCIneoP and pCIneoL(c) (or pCIneo as a negative control).

Plasmid ratio	NP, P & L	NP, P & pCIneo	
pOLTV553N0	$3.1 \times 10^4$	$2.7 \times 10^3$	11.7
pOLTV553N1	4.1	5.2	7.9
pOLTV553N2	3.1	3.1	10.0
pOLTV553N3	35.9	3.6	100.8
pOLTV553N4	1.9	4.6	4.1
pOLTV553N5	1.0	4.1	2.5